

## Catalogue of inland fishes of Chanthaburi Province, Eastern Gulf of Thailand Drainages

Sitthi Kulabtong<sup>1</sup>, Yannan Soonthornkit<sup>2</sup> & Nipaporn Churaroum<sup>2</sup>

<sup>1</sup>Save wildlife of Thailand, Wangnoi District, Ayutthaya Province, Thailand

<sup>2</sup>Fisheries Program, Faculty of Agro-Industrial Technology, Rajamangala University of Technology Tawan-ok Chantaburi Campus Chantaburi, Thailand

\*Corresponding author, e-mail: kulabtong2011@hotmail.com

### ABSTRACT

The present paper reports on a catalogue of inland fishes in Chanthaburi Province, Eastern Gulf of Thailand Drainages. All the species encountered in this region, belonging to 18 orders, 73 families and 229 species, are listed. In particular, *Crossocheilus reticulatus* (Fowler, 1934) (Cypriniformes Cyprinidae), *Pangio anguillaris* (Vaillant, 1902) and *P. oblonga* (Valenciennes, 1846) (Cypriniformes Cobitidae) are new records for Chanthaburi Province, entered in December 2007 and February 2012; description and distribution data of the three new records are provided here.

### KEY WORDS

*Crossocheilus reticulatus*; *Pangio anguillaris*; *Pangio oblonga*; Chanthaburi; Thailand.

Received 09.07.2015; accepted 22.08.2015; printed 30.09.2015

### INTRODUCTION

The Asian Chanthaburi Basin originates at Kit-chakut-Soidao mountain range. This river system runs through Chanthaburi Province, East Thailand, and flows into the Upper Gulf of Thailand at Laem Sing estuary, with a total length of about 123 kilometers. Chanthaburi Basin is a very important river basin but in some areas of it, such as Pongnum Ron District and South Soidao mountain range, very little is known about inland fishes population. At the present moment available data are scarce and fragmented making it difficult to use them.

A survey project aimed at studying freshwater fishes in Chanthaburi Province (see Fig. 1) was carried out in December 2007 and February 2012. We separated this area into two: a pool and a small stream in the mountain; te pool showed transparent slowly waters with an average width of about 20 m,

average depth less than 1 m, and a combination of clay and sand on the bottom; the stream had transparent and running fast waters with an average width of about 15 m, average depth ranging from 0.3–1.0 m, and a combination of rock and sand on the bottom.

We found specimens of *Crossocheilus reticulatus* (Fowler, 1934) (Cypriniformes Cyprinidae), *Pangio anguillaris* (Vaillant, 1902) and *P. oblonga* (Valenciennes, 1846) (Cypriniformes Cobitidae) in Chanthaburi Province. Three species are new records in this area and are reported for the first time in this paper; for previous reviews, see Fowler, 1934; Smith, 1945; Sontirat, 1976; Suvatti, 1981; Karnasuta, 1993; Kottelat & Lim, 1993; Monkolprasit et al., 1997; Robert, 1997; Robert, 1998; Ng & Kottelat, 2000; Soonthornkit, 2001; Sontirat et al., 2006).

ABBREVIATIONS. Standard length = SL; head length = HL.

### RESULTS




Figure 1. Study area: Chanthaburi Province, East Thailand.

### CATALOGUE OF INLAND FISHES IN CHANTHABURI PROVINCE, EAST THAILAND

According to known literature (see above) and present study, inland fishes in Chanthaburi Province belong to 18 orders, 73 families and 229 species.

Order ORECTOLOBIFORMES Applegate, 1972  
Family HEMISCYLLIIDAE Gill, 1862

*Chiloscyllium indicum* (Gmelin, 1789)  
*Chiloscyllium plagiosum* (Anonymous [Bennett], 1830)

Order RAJIFORMES Berg, 1940  
Family GYMNURIDAE Fowler, 1934

*Gymnura micrura* (Bloch et Schneider, 1801)

Order OSTEOGLOSSIFORMES Berg, 1940

Family OSTEOGLOSSIDAE Bonaparte, 1832

*Scleropages formosus* (Müller et Schlegel, 1844)

Family NOTOPTERIDAE Bleeker, 1859

*Notopterus notopterus* (Pallas, 1780)

Order CLUPEIFORMES Bleeker, 1959

Family CLUPEIDAE Cuvier, 1817

*Anodontostoma chacunda* (Hamilton, 1822)

*Clupeichthys goniognathus* Bleeker, 1855

*Sardinella gibbosa* (Bleeker, 1849)

Family ENGRAULIDAE Gill, 1861

*Stolephorus indicus* (van Hasselt, 1823)

*Stolephorus insularis* Hardenberg, 1933

*Stolephorus tri* (Bleeker, 1852)

*Stolephorus waitei* Jordan et Seale, 1926

*Thryssa hamiltonii* Gray, 1835

*Thryssa mystax* (Bloch et Schneider, 1801)

*Thryssa setirostris* (Broussonet, 1782)

Order CYPRINIFORMES Bleeker, 1859

Family CYPRINIDAE Swainson, 1839

*Barbodes rhombeus* Kottelat, 2000

*Barbodes aurotaeniatus* (Tirant, 1885)

*Barbonymus gonionotus* (Bleeker, 1850)

*Barbonymus schwanenfeldii* (Bleeker, 1853)

*Barilius koratensis* (Smith, 1931)

*Cirrhinus microlepis* Sauvage, 1878

*Crossocheilus reticulatus* (Fowler, 1934)

*Cyclocheilichthys apogon* (Valenciennes, 1842)

*Danio albolineatus* (Blyth, 1860)

*Esomus metallicus* Ahl, 1923

*Garra fasciacauda* Fowler, 1937

*Garra cambodgiensis* (Tirant, 1883)

*Hampala macrolepidota* Kuhl et Van Hasselt, 1823

*Hypsibarbus vernayi* (Norman, 1925)

*Labiobarbus leptochela* (Valenciennes, 1842)

*Labiobarbus lineatus* (Sauvage, 1878)

*Laubuca siamensis* Fowler, 1939

*Mystacoleucus marginatus* (Valenciennes, 1842)

*Neolissochilus sumatrana* (Weber et de Beaufort, 1916)

*Neolissochilus stracheyi* (Day, 1871)

- Oreichthys parvus* Smith, 1933  
*Osteochilus microcephalus* (Valenciennes, 1842)  
*Osteochilus schlegelii* (Bleeker, 1851)  
*Osteochilus waandersii* (Bleeker, 1852)  
*Osteochilus vittatus* (Valenciennes, 1842)  
*Osteochilus lini* Fowler, 1935  
*Oxygaster anomalura* Van Hasselt, 1823  
*Parachela maculicauda* (Smith, 1934)  
*Parachela siamensis* (Günther, 1868)  
*Poropuntius bantamensis* (Rendahl, 1920)  
*Poropuntius deauratus* (Valenciennes, 1842)  
*Poropuntius normani* Smith, 1931  
*Puntius leiacanthus* (Bleeker, 1860)  
*Puntius masyai* Smith, 1945  
*Puntigrus partipentazona* (Fowler, 1934)  
*Puntius brevis* (Bleeker, 1850)  
*Rasbora dusonensis* (Bleeker, 1851)  
*Rasbora trilineata* Steindachner, 1870  
*Rasbora borapetensis* Smith, 1934  
*Rasbora myersi* Brittan, 1954  
*Rasbora paviana* Tirant, 1885  
*Systemus orphoides* (Valenciennes, 1842)  
*Trigonostigma heteromorpha* (Duncker, 1904)
- Family GYRINOCHEILIDAE T.N. Gill, 1905  
*Gyrinocheilus aymonieri* (Tirant, 1883)
- Family COBITIDAE Swainson, 1838  
*Acanthopsoides gracilentus* (Smith, 1945)  
*Acantopsis choirorhynchos* (Bleeker, 1854)  
*Barbus diabolica* Roberts, 1989  
*Lepidocephalichthys hasselti* (Valenciennes, 1846)  
*Lepidocephalichthys berdmorei* (Blyth, 18609  
*Pangio anguillaris* (Vaillant, 1902)  
*Pangio kuhlii* (Valenciennes, 1846)  
*Pangio oblonga* (Valenciennes, 1846)  
*Serpenticobitis zonata* Kottelat, 1998  
*Yasuhikotakia sidthimunki* (Klausewitz, 1959)
- Family BALITORIDAE Swainson, 1839  
*Balitora annamitica* Kottelat, 1988  
*Homaloptera modesta* (Vinciguerra, 1890)  
*Homaloptera sexmaculata* Fowler, 1934  
*Homaloptera orthogoniata* Vaillant, 1902  
*Homaloptera smithi* Hora, 1932  
*Homaloptera zollingeri* Bleeker, 1853  
*Nemacheilus binotatus* Smith, 1933  
*Nemacheilus masyai* Smith, 1933
- Nemacheilus pallidus* Kottelat, 1990  
*Nemacheilus platiceps* Kottelat, 1990  
*Schistura deignani* (Smith, 1945)  
*Schistura kohchangensis* (Smith, 1933)  
*Schistura nicholsi* (Smith, 1933)  
*Schistura schultzi* (Smith, 1945)  
*Schistura sexcauda* (Fowler, 1937)
- Order SILURIFORMES Cuvier, 1816  
Family AMBLYCIPITIDAE Day, 1873  
*Amblyceps foratum* Ng et Kottelat, 2000
- Family AKYSIDAE Gill, 1861  
*Akysis maculipinnis* Fowler, 1934  
*Pseudobagarius macronemus* (Bleeker, 1860)
- Family SISORIDAE Bleeker, 1858  
*Glyptothorax fuscus* Fowler, 1934  
*Glyptothorax major* (Boulenger, 1894)  
*Glyptothorax trilineatus* Blyth, 1860
- Family SILURIDAE Rafinesque, 1815  
*Ompok siluroides* Lacepède, 1803  
*Pterocryptis torrentis* (Kobayakawa, 1989)  
*Silurichthys phaiosoma* (Bleeker, 1851)  
*Silurichthys hasseltii* Bleeker, 1858  
*Silurichthys schneideri* Volz, 1904
- Family PLOTOSIDAE Bleeker, 1858  
*Plotosus canius* Hamilton, 1822
- Family CLARIIDAE Bonaparte, 1845  
*Clarias batrachus* (Linnaeus, 1758)
- Family BAGRIDAE Bleeker, 1858  
*Hemibagrus nemurus* (Valenciennes, 1840)  
*Leiocassis micropogon* (Bleeker, 1852)  
*Mystus cavasius* (Hamilton, 1822)  
*Mystus castaneus* Ng, 2002  
*Pseudomystus stenomus* (Valenciennes, 1840)  
*Pseudomystus siamensis* (Regan, 1913)

## Family ARIIDAE Bleeker, 1858

- Arius maculatus* (Thunberg, 1792)  
*Arius venosus* Valenciennes, 1840  
*Batrachocephalus mino* (Hamilton, 1822)  
*Hexanemichthys sagor* (Hamilton, 1822)  
*Osteogeneiosus militaris* (Linnaeus, 1758)  
*Plicofollis argyropleuron* (Valenciennes, 1840)

## Order AULOPIFORMES D.E. Rosen, 1973

## Family SYNODONTIDAE Gill, 1861

- Saurida tumbil* (Bloch, 1795)

## Order GADIFORMES Goodrich, 1909

## Family BREGMACEROTIDAE Gill, 1872

- Bregmaceros mcclellandi* Thompson, 1840

## Order BATRACHOIDIFORMES Goodrich, 1909

## Family BATRACHOIDIDAE Jordan, 1896

- Allenbatrachus grunniens* (Linnaeus, 1758)

## Order ATHERINIFORMES D.E. Rosen, 1966

## Family PHALLOSTETHIDAE Regan, 1916

- Neostethus bicornis* Regan, 1916  
*Neostethus lankesteri* Regan, 1916  
*Phenacostethus smithi* Myers, 1928

## Order BELONIFORMES L.S. Berg, 1937

## Family BELONIDAE Bonaparte, 1835

- Xenentodon cancila* (Hamilton, 1822)

## Family EXOCOETIDAE Risso, 1827

- Cheilopogon arcticeps* (Günther, 1866)

## Family HEMIRAMPHIDAE Gill, 1859

- Dermogenys siamensis* Fowler, 1934  
*Zenarchopterus ectuntio* (Hamilton, 1822)

## Order SYNGNATHIFORMES Berg, 1940

## Family CENTRISCIDAE Bonaparte, 1831

- Centriscus scutatus* Linnaeus, 1758

## Family SYNGNATHIDAE Bonaparte, 1831

- Doryichthys boaja* (Bleeker, 1851)  
*Doryichthys martensi* (Peters, 1868)  
*Hippocampus kuda* Bleeker, 1852

## Order CYPRINODONTIFORMES L.S. Berg, 1940

## Family APLOCHEILIDAE Bleeker, 1859

- Aplocheilus panchax* (Hamilton, 1822)

## Order SYNBRANCHIFORMES Nelson, 1994

## Family SYNBRANCHIDAE Bonaparte, 1835

- Monopterus albus* (Zuiw, 1793)

## Family MASTACEMBELIDAE Swainson, 1839

- Macrognathus circumcinctus* (Hora, 1924)  
*Macrognathus siamensis* (Günther, 1861)  
*Mastacembelus armatus* (Lacepède, 1800)  
*Mastacembelus favus* Hora, 1924

## Order SCORPAENIFORMES Greenwood et al., 1966

## Family APISTIDAE Gill, 1859

- Apistus carinatus* (Bloch et Schneider, 1801)

## Family PLATYCEPHALIDAE Swainson, 1839

- Grammoplites scaber* (Linnaeus, 1758)  
*Platycephalus indicus* (Linnaeus, 1758)

## Family SYNANCEIIDAE Swainson, 1839

- Inimicus didactylus* (Pallas, 1769)  
*Trachicephalus uranoscopus* (Bloch et Schneider, 1801)

## Family TETRAROGIDAE Smith, 1949

- Paracentropogon longispinis* (Cuvier, 1829)  
*Pterois russelli* Bennett, 1831

- Order PERCIFORMES Bleeker, 1859  
 Family AMBASSIDAE Klunzinger, 1870  
*Parambassis siamensis* (Fowler, 1937)
- Family LATIDAE Jordan, 1888  
*Lates calcarifer* (Bloch, 1790)
- Family APOGONIDAE Günther, 1859  
*Apogon hyalosoma* Bleeker, 1852  
*Apogon multitaeniatus* Cuvier, 1828
- Family SILLAGINIDAE Richardson, 1846  
*Sillago sihama* (Forsskål, 1775)
- Family CARANGIDAE Rafinesque, 1815  
*Alectis ciliaris* (Bloch, 1787)  
*Carangoides praeustus* (Anonymous [Bennett], 1830)  
*Caranx sexfasciatus* Quoy et Gaimard, 1825  
*Parastromateus niger* (Bloch, 1795)  
*Scomberoides lysan* (Forsskål, 1775)
- Family LEIOGNATHIDAE Gill, 1893  
*Gazza minuta* (Bloch, 1795)  
*Leiognathus daura* (Cuvier, 1829)  
*Secutor ruconius* (Hamilton, 1822)
- Family LUTJANIDAE Gill, 1861  
*Lutjanus sanguineus* (Cuvier, 1828)  
*Lutjanus vitta* (Quoy et Gaimard, 1824)
- Family LOBOTIDAE Gill, 1861  
*Lobotes surinamensis* (Bloch, 1790)
- Family GERREIDAE Bleeker, 1859  
*Gerres erythrourus* (Bloch, 1791)  
*Gerres setifer* (Hamilton, 1822)
- Family HAEMULIDAE Gill, 1885  
*Plectorhinchus nigrus* (Cuvier, 1830)
- Pomadasys hasta* (Bloch, 1790)  
*Pomadasys maculatus* (Bloch, 1793)
- Family SPARIDAE Rafinesque, 1818  
*Acanthopagrus berda* (Forsskål, 1775)
- Family POLYNEMIDAE Rafinesque, 1815  
*Eleutheronema tetradactylum* (Shaw, 1804)  
*Eleutheronema tridactylum* (Bleeker, 1849)
- Family SCIAENIDAE Cuvier, 1829  
*Nibea soldado* (Lacepède, 1802)  
*Pennahia argentata* (Houttuyn, 1782)
- Family MULLIDAE Rafinesque, 1815  
*Upeneus sulphureus* Cuvier, 1829  
*Upeneus tragula* Richardson, 1846  
*Upeneus vittatus* (Forsskål, 1775)
- Family MONODACTYLIDAE Jordan et Evermann, 1898  
*Monodactylus argenteus* (Linnaeus, 1758)
- Family TAXOTIDAE Bleeker, 1859  
*Toxotes jaculatorius* (Pallas, 1767)
- Family POMACENTRIDAE Bonaparte, 1831  
*Abudefduf bengalensis* (Bloch, 1787)  
*Neopomacentrus taeniurus* (Bleeker, 1856)  
*Pristotis obtusirostris* (Günther, 1862)
- Family NANDIDAE Bleeker, 1852  
*Nandus nebulosus* (Gray, 1835)  
*Pristolepis fasciata* (Bleeker, 1851)
- Family TERAPONTIDAE Richardson, 1842  
*Pelates sexlineatus* (Quoy et Gaimard, 1825)  
*Terapon theraps* Cuvier, 1829
- Family CEPOLIDAE Rafinesque, 1815  
*Acanthocepola limbata* (Valenciennes, 1835)

## Family LABRIDAE Cuvier, 1816

*Halichoeres argus* (Bloch et Schneider, 1801)  
*Halichoeres nigrescens* (Bloch et Schneider, 1801)  
*Thalassoma lunare* (Linnaeus, 1758)

## Family BLENNIIDAE Rafinesque, 1810

*Istiblennius dussumieri* (Valenciennes, 1836)  
*Petroscirtes mitratus* Rüppell, 1830

## Family CALLIONYMIDAE Bonaparte, 1831

*Dactylopus dactylopus* (Valenciennes, 1837)

## Family ELEOTRIDAE Bonaparte, 1835

*Butis butis* (Hamilton, 1822)  
*Butis gymnopomus* (Bleeker, 1853)  
*Ophiocara porocephala* (Valenciennes, 1837)

## Family GOBIIDAE Cuvier, 1816

*Acentrogobius bontii* (Bleeker, 1849)  
*Acentrogobius caninus* (Valenciennes, 1837)  
*Acentrogobius chlorostigmatooides* (Bleeker, 1849)  
*Acentrogobius viridipunctatus* (Valenciennes, 1837)  
*Callogobius hasseltii* (Bleeker, 1851)  
*Cryptocentrus callopterus* Smith, 1945  
*Cryptocentrus diproctaenia* Bleeker, 1876  
*Eugnathogobius oligactis* (Bleeker, 1875)  
*Glossogobius giuris* (Hamilton, 1822)  
*Mahidolia mystacina* (Valenciennes, 1837)  
*Oxyurichthys microlepis* (Bleeker, 1849)  
*Parapocryptes serperaster* (Richardson, 1846)  
*Periophthalmus barbarus* (Linnaeus, 1766)  
*Priolepis semidoliata* (Valenciennes, 1837)  
*Scartelaos histophorus* (Valenciennes, 1837)  
*Valenciennea muralis* (Valenciennes, 1837)  
*Yongeichthys nebulosus* (Forsskål, 1775)

## Family SCATOPHAGIDAE Gill, 1883

*Scatophagus argus* (Linnaeus, 1766)

## Family SPHYRAENIDAE Rafinesque, 1815

*Sphyraena barracuda* (Edwards, 1771)  
*Sphyraena flavicauda* Rüppell, 1838  
*Sphyraena obtusata* Cuvier, 1829

## Family SCOMBRIDAE Rafinesque, 1815

*Scomberomorus commerson* (Lacepède, 1800)  
*Scomberomorus guttatus* (Bloch et Schneider, 1801)

## Family DREPANEIDAE Gill, 1872

*Drepane punctata* (Linnaeus, 1758)

## Family ANABANTIDAE Bonaparte, 1831

*Anabas testudineus* (Bloch, 1792)

## Family OSPHRONEMIDAE van der Hoeven, 1832

*Betta taeniata* Regan, 1910  
*Betta prima* Kottelat, 1994  
*Trichogaster trichopterus* (Pallas, 1770)  
*Trichopsis pumila* (Arnold, 1936)  
*Trichopsis vittata* (Cuvier, 1831)

## Family CHANNIDAE Fowler, 1934

*Channa gachua* (Hamilton, 1822)  
*Channa lucius* (Cuvier, 1831)  
*Channa striata* (Bloch, 1793)

## Family CYNOGLOSSIDAE Jordan, 1888

*Cynoglossus lingua* Hamilton, 1822

Order PLEURONECTIFORMES Linnaeus, 1758  
Family SOLEIDAE Bonaparte, 1833

*Brachirus orientalis* (Bloch et Schneider, 1801)

## Family PARALICHTHYIDAE Regan, 1910

*Pseudorhombus arsius* (Hamilton, 1822)  
*Pseudorhombus oligodon* (Bleeker, 1854)

Order TETRAODONTIFORMES L.S. Berg, 1940  
Family BALISTIDAE Rafinesque, 1810

*Abalistes stellatus* (Anonymous, 1798)

## Family DIODONTIDAE Bonaparte, 1835

*Diodon holocanthus* Linnaeus, 1758

## Family TETRAODONTIDAE Bonaparte, 1831

- Arothron leopardus* (Day, 1878)  
*Arothron stellatus* (Bloch et Schneider, 1801)  
*Tetraodon cambodgiensis* Chabanaud, 1923  
*Tetraodon fluviatilis* Hamilton, 1822  
*Tetraodon leiurus* Bleeker, 1851

## Family TRIACANTHIDAE Bleeker, 1859

- Triacanthus biaculeatus* (Bloch, 1786)

## NEW RECORDS FOR CHANTHABURI PROVINCE

*Crossocheilus reticulatus* (Fowler, 1934)

EXAMINED MATERIAL. 3 specimens, 48.3–61.1 mm SL, Pongnum Ron District, Chanthaburi Province, East Thailand, XII.2007 and II.2012, legit S. Kulabtong (Fig. 2).

DESCRIPTION. *Crossocheilus reticulatus* is compress, body depth is 26.23–27.1 %SL. Body width is 12.28–12.35 %SL. Scales in lateral series are medium to large, lateral series scales are 31–33. Head length is 23.52–23.75 %SL. The eyes is large, eye diameter is 36.7–37.1 %HL. Snout length is long, with 37.41–38.2 %HL and interorbital width is 40.29–40.46 % HL. Dorsal fin origin is anterior pelvic fin origin, predorsal fin length is 48.22–49.31 %SL, prepectoral fin length is 22.87–23.69 %SL, prepelvic fin length is 51.8–53.47 %SL and preanal fin length is 75.63–78.24 %SL. Caudal peduncle depth is 11.2–12.52 %SL. Pectoral fin is short not reaching beyond anus, the pectoral fin is 18.2–20.98 %SL long with 13–14 branched fin rays. Pelvic fin is short not reaching beyond anus, the pelvic fin is 18.95–20.33 %SL long with 7–8 branched fin rays. Anal fin base is shorter than dorsal fin base, the anal fin base length is 8.8–10.6 %SL, dorsal fin with 2 unbranched rays and 10–11 branched rays and dorsal fin base length is 16.07–17.4 %SL.

DISTRIBUTION. This species is known from peninsular Thailand, Mekong Basin in Indochina, Chao Phraya Basin and Maeklong Basin, Thailand. New record for Chanthaburi Province.

*Pangio anguillaris* (Vaillant, 1902)

EXAMINED MATERIAL. 1 specimen, 60 mm SL, from Pongnum Ron District, Chanthaburi Province, East Thailand, XII.2007, legit S. Kulabtong (Fig. 3).

DESCRIPTION. *Pangio anguillaris* is compress, body depth is 6.05 %SL. Body width is 1.96 %SL. Head length is 10.46 %SL. The eyes are small, eye diameter is 1.56 %HL. Snout length is long, with 34.38 %HL and interorbital width is 32.6 % HL. Dorsal fin origin is posterior pelvic fin origin, predorsal fin length is 67.97 %SL, prepectoral fin length is 9.48 %SL, prepelvic fin length is 50.33 %SL and preanal fin length is 69.12 %SL. Caudal peduncle depth is 3.76 %SL. Pectoral fin is short not reaching beyond anus, the pectoral fin is 6.86 %SL long with 5 branched fin rays. Pelvic fin is short not reaching beyond anus, the pelvic fin is 6.54 %SL long with 6 branched fin rays. Anal fin base is shorter than dorsal fin base, the anal fin base




Figure 2. *Crossocheilus reticulatus*, 57.6 mm SL. Figure 3. *Pangio anguillaris*, 60.8 mm SL. Figure 4. *Pangio oblonga*, 45.6 mm SL.

length is 0.16 %SL, dorsal fin shows 1 unbranched and 7 branched rays; dorsal fin base length is 0.16 %SL.

**DISTRIBUTION.** This species is known from Malaysia, Indonesia, peninsular Thailand, Mekong Basin in Indochina, Chao Phraya Basin and Rayong River, Thailand. New record for Chanthaburi Province.

### *Pangio oblonga* (Valenciennes, 1846)

**EXAMINED MATERIAL.** One specimen, 43.1 mm SL, from Pongnum Ron District, Chanthaburi Province, East Thailand, XII.2007, legit S. Kulabtong (Fig. 4).

**DESCRIPTION.** *Pangio oblonga* is compress, body depth is 11.03 %SL. Body width is 5.75 %SL. Head length is 19.08 %SL. The eyes are small, eye diameter is 15.66 %HL. Snout length is long, with 40.96 %HL and interorbital width is 30.12 % HL. Dorsal fin origin is posterior pelvic fin origin, pre-dorsal fin length is 63.68 %SL, prepectoral fin length is 20.0 %SL, prepelvic fin length is 59.31 %SL and preanal fin length is 82.53 %SL. Caudal peduncle depth is 6.44 %SL. Pectoral fin is short not reaching beyond anus, the pectoral fin is 10.11 %SL long with 5 branched fin rays. Pelvic fin is short not reaching beyond anus, the pelvic fin is 6.21 %SL long with 6 branched fin rays. Anal fin base is short than dorsal fin base, the anal fin base length is 0.23 %SL, dorsal fin with 1 unbranched and 7 branched rays; dorsal fin base length is 0.23 %SL.

**DISTRIBUTION.** This species is known from Malaysia, Indonesia, peninsular Thailand, Mekong Basin in Indochina, Chao Phraya Basin and Rayong River, Thailand. New record for Chanthaburi Province.

### ACKNOWLEDGMENTS

The authors are grateful to anonymous referees for reviewing this manuscript and a special thank to all partners for helping in collecting the specimens employed in this study.

### REFERENCES

- Fowler H.W., 1934. Zoological results of the third De Schauensee Siamese Expedition, Part I. Proceedings of the Academy of Natural Sciences of Philadelphia, 86: 67–163.
- Karnasuta J., 1993. Systematic revision of Southeastern Asiatic cyprinid fish genus *Osteochilus* with description of two new species and a new subspecies. Kasetsart University Fishery Research Bulletin, 19: 105 pp.
- Kottelat M. & Lim K.K.P., 1993. A review of the eel-loaches of the genus *Pangio* (Teleostei: Cobitidae) from the Malay Peninsula, with descriptions of six new species. Raffles bulletin of zoology, 41: 203–249.
- Monkolprasit S., Sontirat S., Vimolohakarm S. & Songsirikul T., 1997. Checklist of fishes in Thailand. Office of environmental policy and planning, Thailand.
- Ng H.H. & Kottelat M., 2000. A review of the genus *Amblyceps* (Osteichthyes: Amblycipitidae) in Indochina, with descriptions of five new species. Ichthyological Exploration of Freshwaters, 11: 335 – 348.
- Roberts T.R., 1997. Systematic revision of the tropical Asian labeoin cyprinid fish genus *Cirrhinus*, with descriptions of new species and biological observations on *C. lobatus*. Natural History Bulletin of the Siam Society, 45: 171200, 505203.
- Roberts T.R., 1998. Freshwater fugu or pufferfishes of the genus *Tetraodon* from the Mekong basin, with descriptions of two new species. Ichthyological Research, 45: 225200, 505234.
- Smith H.M., 1945. The fresh-water fishes of Siam, or Thailand. National Museum Bulletin 188. U.S. Government Printing office, Washington D.C. 622.
- Sontirat S., 1976. Revision of the Southeastern Asiatic cyprinid fish Genus *Cyclocheilichthys*, Ph.D. Thesis, University of Michigan, Ann. Arbor, 133 pp.
- Sontirat S., Tunchareon S. & Soothornkit Y., 2006. Fish species diversity in the areas of national parks and wildlife sanctuaries in the five eastern provinces of Thailand. The 44th proceeding of the Kasetsart University conference, pp. 60 – 67.
- Soothornkit Y., 2001. Fish species diversity in Khao Khitchakut National Parks and Khao Soi Dao wildlife sanctuaries. Master Thesis, Kasetsart University, Thailand, 298 pp.
- Suvatti C., 1981. Fishes of Thailand. Royal Institute, Thailand, 379 pp.