# Terrestrial gastropods (Mollusca Gastropoda) from Lepini Mountains (Latium, Italy): a first contribution

Alessandro Hallgass<sup>1</sup> & Angelo Vannozzi<sup>2</sup>

<sup>1</sup>Via della Divina Provvidenza 16, 00166 Rome, Italy; e-mail: hallgass@hotmail.com <sup>2</sup>Via M.L. Longo 8, 00151 Rome, Italy; e-mail: ang.vannozzi@gmail.com

#### ABSTRACT

Lepini Mountains are a calcareous massif that forms the southern pre-Apennines of Latium (Italy), reaching a maximum altitude of 1536 m. Notwithstanding their central position and the low height reached, the malacofauna of Lepini Mountains has been long neglected and species composition was never reported so far. In this contribution, a preliminary investigation of the terrestrial gastropods (Mollusca Gastropoda) occurring in the Lepini Mountains is reported. At least 43 species are recorded. Several species already reported from Central Apennines occur. The most remarkable findings include a hitherto unrecorded population of *Medora* sp. (Clausiliidae) and the occurrence of two distinct forms ascribable to *Jaminia quadridens* s.1.

**KEY WORDS** Terrestrial gastropods; biodiversity; Lepini Mountains; Italy.

Received 28.02.2016; accepted 20.03.2016; printed 30.03.2016 Proceedings of the Ninth Malacological Pontine Meeting, October 3rd-4th, 2015 - San Felice Circeo, Italy

## INTRODUCTION

Lepini Mountains, together with Aurunci and Ausoni Mountains, form the southern pre-Apennines of Latium (Italy). They are positioned about 50 km SE of Rome and extend in NE-SW direction (Fig. 1). They are separated from Central Apennines by the Sacco Valley and face the Pontine alluvial plain in the south. Lepini Mountains are comprised of two parallel chains directed in NE-SW direction, separated by the deep Montelanico-Carpineto-Maenza tectonic line. The western chain is comprised by Mount Lupone (1378 m) and Mount Semprevisa group (1536 m, the highest peak), whereas the eastern chain quickly slopes down to the Sacco Valley and is comprised of Mount Gemma, Mount Malaina, Mount S. Marino and Mount Alto, all of which reach heights around 1400 m.

Lepini Mountains are mainly comprised of limestone of Cretaceous age (Sani et al., 2004). The whole massif shows to intense karst phenomena. As a consequence, in the Lepini Mountains no permanent water body occurs, whereas several springs appear at the base of the massif, the best known of which gives rise to the Oasis of Ninfa (Amori et al., 2002).

The vegetation is mainly comprised of holm oak, chestnut and mixed woods at medium-low altitudes and beech forests at medium-high altitudes. Large portions of territory are occupied by grassland mainly used as pasture.

The invertebrate fauna of Lepini Mountains have been studied in some detail with regard to arthropods (Corsetti et al., 2015). Several studies focused on hypogean fauna (Sbordoni, 1971; Latella, 1995; Nardi et al., 2002). Moreover, some endemisms have been reported (Sbordoni, 1971;



Figure 1. Studied area: Lepini Mountains (southern Latium, Italy). Numbers indicate the sampled stations listed in Table 1.

Stn.	Locality	Coordinates	Alt. (m)	Environment
1	Pass to Campo di Segni	41.672649° N, 12.987930° E	1015	Pasture with stones
2	After pass to Campo di Segni	41.670720° N, 12.993717° E	960	Rocks with low vegetation
3	Close to Campo di Segni	41.667926° N, 12.990029° E	880	Pasture and bushes wit stones
4	Mount Erdigheta	41.56410° N, 13.119629° E	1046	Beech forest with rocks
5	Mount Erdigheta	41.562092° N, 13.120613° E	1115	Pasture with stones
6	Mount Semprevisa	41.572228° N, 13.092678° E	1250– 1400	Beech forest with rocks
7	Mount Semprevisa, top	41.571147° N, 13.091063° E	1490	Stones on the top
8	Carpineto, Pian della Faggeta	41.575702° N, 13.103665° E	930	Rocks with low vegetation and residuary beeches
9	Bassiano, road to Sempre- visa, near the spring	41.552975° N, 13.047529° E	590	Holm oak wood with rocks
10	Bassiano, road to Semprevisa	41.558473° N, 13.059121° E	864	Clearing in holm oak wood
11	Campo Rosello	41.563711° N, 13.077542° E	1174	Pasture with stones
12	Campo Rosello	41.571987° N, 13.072649° E – 41.574551° N, 13.074571° E	1250– 1410	Pasture with stones

Table 1. List of the stations: Lepini Mountains (southern Latium, Italy).

Pace, 1975; Magrini, 2005). However, the molluscan fauna of Lepini Mountains was never studied so far. Only four species occurring in hypogean environments were reported (*Discus rotundatus*, *Campylaea planospira*, *Daudebardia brevipes* and *Oxychilus draparnaudi*), none of which strictly hypogean (Latella, 1995).

In this contribution, we report the results of a first survey aimed at assessing the biodiversity of terrestrial gastropods of Lepini Mountains.

#### MATERIAL AND METHODS

A total of 12 stations along the western chain have been sampled between April and June 2015 (see Table 1). Sampling was carried out only in natural environments. As a consequence, species recorded only from urban areas, such as Cornu aspersum (O.F. Müller, 1774) found in the town of Bassiano, were ruled out. Additionally, freshwater or hypogean environments were not considered. For the nomenclature, we mainly referred to the checklist of the species of the Italian fauna (Bodon et al., 1995; Manganelli et al., 1995, 1998, 2000). For the suprageneric nomenclature, we referred to Bouchet & Rocroi (2005). All specimens here illustrated were collected from the Lepini Mountains. Shell length and width were measured parallel and perpendicular to the axis of the shell, respectively, with calipers to the nearest 0.1 mm.

## **RESULTS AND DISCUSSION**

At least 44 species of terrestrial gastropods occur in the Lepini Mountains (see Table 2). The most speciose family is the Clausiliidae, with 7 recorded species. The clausiliid *Leucostigma candidescens* is by far the commonest and widespread species, occurring in almost all calcareous outcrops, either exposed or shaded, often associated with other calciophilous species such as *Cochlostoma* cf. *adamii*, *Marmorana signata*, *Granaria apennina* and *Medora* sp. We agree with Fehér et al. (2010) who indicate th species of the genus *Granaria* Held, 1838 occurring in the Italian peninsula as *G. apennina*. *Medora* sp. was found only in a single station inside the beech forest of Mount Semprevisa. The unexpected finding of this population confirms that the current knowledge of the genus Medora H. et A. Adams, 1855 in Italy is far from being exhaustive (Giusti et al., 1986; Nordsieck, 2012; Colomba et al., 2012). Cochlodina laminata and C. bidens occur in sympatry in the beech forest. According to Opinion 2355, the Apennine species so far known as Cochlodina incisa (Küster, 1876) should be indicated as C. bidens (Linnaeus, 1758) (Kadolsky, 2009; ICZN, 2015). They are readily distinguished by the development of palatal plicae. In fact, while in the former palatal plicae are truncated at the level of the clausilium, in the latter both the principal and the lower palatal plicae prolong internally. Moreover, an additional intermediate palatal plica often occurs in the latter. Cochlodina bidens in Lepini Mountains shows a stout shell also found in specimens from other localities of Latium, such as the holm oak woods of Mount Circeo and Macchia Grande (Fiumicino) (Hallgass & Vannozzi, 2014).

The populations of Cochlostoma cf. adamii have been studied by Zallot et al. (2015) in the generic revision of the family Cochlostomatidae and assigned to the subgenus Turritus Westerlund, 1883. Cochlostoma adamii group is comprised of several forms reported with different nominal taxa occurring from Central Apennines to Sicily, whose taxonomy needs to be clarified. The marquise Paulucci (1881) noted the occurrence of forms close to Pomatias adamii in the Central Apennines and described "Pomatias adamii Var. Carseolanus" from Carsoli (Abruzzi). Cochlostoma cf. adamii from Lepini Mountains is different from C. cassiniacum (Saint Simon in Paulucci, 1878) from Cassino and Sterrone (both Latium), though belonging to the same subgenus Turritus (Zallot, comm. pers.).

On the whole, the beech forest shows the greatest biodiversity, with 28 recorded species. Among them, there are several species commonly found in beech forests of Central Apennines (Giusti et al., 1985). However, a few of them deserve some comments. *Acicula* sp. was recorded from a worn fragment. The closest finding of this genus is *A. szigethyannae* Subai, 1977 from Val d'Arano (Ovindoli, Abruzzi). Conversely, *Platyla similis* is recorded from several localities of the Italian peninsula. In particular, it has been reported from the neighbouring Aurunci Mountains (Bodon & Cianfanelli, 2008). *Limax* cf. *maximus* appears with different patterns (Figs. 8 and 9). A completely

COCHLOSTOMATIDAE Cochlostoma cf. adamii (Paulucci, 1879) 4–9, 11, 12 2   ACICULIDAE Acicula sp. 6 11   Platyla similis (Reinhardt, 1880) 6 10   POMTIDAE Pomatise elegans (O.F. Müller, 1774) 2, 3, 9   ORCULIDAE Sphyradium doliolum (Bruguière, 1792) 6, 8   VALLONIIDAE Acamhinula acuileata (O.F. Müller, 1774) 8, 9   ORCULIDAE Sphyradium doliolum (Bruguière, 1792) 6, 8   VALLONIIDAE Acamhinula acuileata (O.F. Müller, 1774) 8, 9   Chondrina avenacea (Bruguière, 1782) 6, 9 14   ENIDAE Jaminia quadriders (O.F. Müller, 1774) (small morphotype) 7, 8 5, 18   ENIDAE Jaminia quadriders (O.F. Müller, 1774) 9 14   ERUSSACIIDAE Cecliolades acciula (O.F. Müller, 1774) 9 15   SUBULINIDAE Rumina decollata (Linnaeus, 1758) 1, 3, 9 14   Cochiodina laminata (Montagu, 1803) 6 23 22   Cochiodina laminata (Montagu, 1801) 6 12 23   Cochiodina laminata (Montagu, 1803)	Family	Species	Stn.	Fig.
COUNDARY IDM Counter and in (Counter, 1877) 14-7, 11, 12 12   ACICULIDAE Acticula sp. 6 11   POMATIIDAE Promatics elegans (O.F. Müller, 1774) 2, 3, 9 0   ORCULIDAE Sphyradium doloidum (Bruguier, 172) 6, 8 0   VALLONIIDAE Acanthinuda aculeata (O.F. Müller, 1774) 8, 9 13   Giinenbergia sororcula (Benoi, 1857) 6 15   Chondrina avenaca (Bruguière, 1792) 6, 9 14   ENIDAE Granaria agennia (Käster, 1850) 4, 8 14   ENIDAE Jaminia quadridens (O.F. Müller, 1774) (small morphotype) 7, 8 5, 18   Jaminia quadridens (O.F. Müller, 1774) 9 14 19   Merdigera obscura (O.F. Müller, 1774) 9 13, 20   CLAUSILIDAE Redora sp. 6 20   Leucostigma candidescers (Rossmäsler, 1835) 1-6, 8, 9, 11, 12 3, 21   Cocholdina laminata (Montagu, 1803) 6 23 22   Sicillaria paestana (Philippi, 1836) 3, 5, 6, 9 14   Macrogastra plicatula (Draparaud, 1801)		Cochlostoma of adamii (Poulucci, 1870)	4 0 11 12	2
ACCOLDAD: <t< td=""><td></td><td><i>Cochiosioma</i> ci. <i>adamii</i> (Faulucci, 1879)</td><td>4-9, 11, 12</td><td>11</td></t<>		<i>Cochiosioma</i> ci. <i>adamii</i> (Faulucci, 1879)	4-9, 11, 12	11
POMATIDAE Pomatise (cennula, 1880) 0 10   POMATIDAE Pomatise legans (O.F. Müller, 1774) 2, 3, 9   ORCULIDAE Sphyradium doliohum (Bruguière, 1792) 6, 8   VALLONIIDAE Acamhinula aculeata (O.F. Müller, 1774) 8, 9 13   Chondrina avenacea (Bruguière, 1792) 6, 8 15   CHONDRINIDAE Granaria apennina (Kister, 1850) 4, 8 14   ENIDAE Trancatellina callicratis (Scacchi, 1833) 9 14   ENIDAE Trancatellina callicratis (Scacchi, 1833) 9 14   Statistic quadriders (O.F. Müller, 1774) (small morphotype) 7, 8 5, 18   Jaminia quadriders (O.F. Müller, 1774) (small morphotype) 4, 5, 7, 8, 12 4, 19   Merdigera obscura (O.F. Müller, 1774) 6, 8 10   SUBULINDAE Reinina decollata (Linnaeus, 1758) 1, 3, 9 13, 20   CLAUSILIIDAE Medora sp. 6 20   Cachiodina binders (Linnaeus, 1758) 6 22   Scillaria paestana (Philippi, 1836) 3, 5, 6, 9 10   Mecrogastra plicatula (Draparnaud, 1801) 6 <td>ACICULIDAE</td> <td>Aciculu Sp.</td> <td>6</td> <td>10</td>	ACICULIDAE	Aciculu Sp.	6	10
POMATIDAE Pomatiae sergatis (O.F. Müller, 17/4) 2, 3, 9   ORCULIDAE Splyradium dolium (Bruguiser, 1792) 6, 8   VALLONIIDAE Acanthinula aculeata (O.F. Müller, 1774) 8, 9 13   Gittenbergia sororcula (Benoi, 1857) 6 15   CHONDRINIDAE Granaria apennina (Küster, 1850) 4, 8   Chondrina avenacea (Bruguièr, 1792) 6, 9 4   VERTIGINIDAE Truncatellina calicratis (Scacchi, 1833) 9 14   ENIDAE Jaminia quadrides (O.F. Müller, 1774) 6, 8 5, 18   Jaminia quadrides (O.F. Müller, 1774) 9 5 5, 18   SUBULINIDAE Ceclibides acicula (O.F. Müller, 1774) 9 6 20   CLAUSLIDAE Reuina decolata (Linnacus, 1758) 1, 3, 9 6 23   Cocholatina laminata (Montagu, 1803) 6 23 6 22   Siciliaria paestana (Philippi, 1836) 3, 5, 6, 9 6 22   Macrogastra plicatula (Draparnaud, 1801) 6 12 10   DISCIDAE Discus rotundatus (O.F. Müller, 1774) 6, 9 12 <td></td> <td>Paratian classers (O.E. Müller, 1774)</td> <td>2 2 0</td> <td>10</td>		Paratian classers (O.E. Müller, 1774)	2 2 0	10
ORCOLIDAESphyraaini achindin (Sugurer, 172)6,8VALLONIIDAEAcamhinia acualicata (O.F. Müller, 1774)8,913Gittenbergia sororcula (Benoit, 1857)615CHONDRINIDAEGranaria apennina (Küster, 1850)4,84,8Chondrina avenacea (Bruguière, 1792)6,99VERTIGINIDAETruncatellina callicratis (Scacchi, 1833)914ENIDAEJaminia quadridens (O.F. Müller, 1774) (small morphotype)7,85,18Jaminia quadridens (O.F. Müller, 1774)6,8914FERUSSACIIDAECecilioides acicula (O.F. Müller, 1774)99SUBULINIDAERumina decollata (Linnaeus, 1758)1,3,91CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1-6,8,9,11,123,21Cochlodina laminata (Montagu, 1803)623Cochlodina laminata (Montagu, 1803)622Siciliaria paestana (Philippi, 1836)3,5,6,91Macrogastra plicatula (Draparmaud, 1801)612DISCIDAEPunctum pygmacum (Draparmaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6,917Vitrea subrinata (Reinhardt, 1871)6,816OXYCHILIDAEVitrea subrinata (Reinhardt, 1871)612DISCIDAEDaudebardia rufa (Draparmaud, 1805)66Oxychilas cf. draparmaud, 1805)6610Oxychilas cf. draparmaud, 1805)61610OXYCHILIDAEVi		Pomanas elegans (O.F. Muller, 1774)	2, 3, 9	
VALLONIDAE Acaminia decueda (D.F. Müller, 1/14) 8, 9 13   Gittenbergia sororcula (Benoit, 1857) 6 15   CHONDRINIDAE Granaria ageminia (Küster, 1850) 4, 8   Chondrina avenacea (Bruguière, 1722) 6, 9 14   ENIDAE Turacatellina callicratis (Scacchi, 1833) 9 14   ENIDAE Jaminia quadridens (O.F. Müller, 1774) (arge morphotype) 7, 8 5, 18   Jaminia quadridens (O.F. Müller, 1774) 9 4, 19 4, 19   Merdigera obscura (O.F. Müller, 1774) 9 5 20   CLAUSLIDAE Cecilioides acicula (O.F. Müller, 1774) 9 6 20   CLAUSLIDAE Rumina decollata (Linnaeus, 1758) 1-6, 8, 9, 11, 12 3, 21   Cocholdina laminata (Montagu, 1803) 6 22 3   Sciciliaria paestana (Philippi, 1836) 3, 5, 6, 9 22 3   Macrogastra plicatula (Draparnaud, 1801) 6 12 21   DISCIDAE Punctum pygmacum (Draparnaud, 1801) 6 12 21   DISCIDAE Discus rotundatus (O.F. Müller, 1774)		Sphyraaium aoitoium (Bruguiere, 1792)	0, 8	10
Childenergia soroccial (Benoti, 1857) 6 13   CHONDRINIDAE Granaria apennina (Küster, 1850) 4, 8   Chondrina avenacea (Bruguière, 1792) 6, 9   VERTIGNIDAE Truncatellina callicratis (Scacchi, 1833) 9   INDAE Jaminia quadridens (O.F. Müller, 1774) (small morphotype) 7, 8 5, 18   Merdigera obscura (O.F. Müller, 1774) (arge morphotype) 4, 5, 7, 8, 12 4, 19   Merdigera obscura (O.F. Müller, 1774) 6, 8 1   SUBULNIDAE Cecitioides acicula (O.F. Müller, 1774) 9 1   CLAUSILIIDAE Meedora sp. 6 20   CLAUSILIIDAE Meedora sp. 6 20   Cochiodina laminata (Montagu, 1803) 6 23   Cochiodina bidens (Linnacus, 1758) 1-6, 8, 9, 11, 12 3, 21   Cochiodina bidens (Linnacus, 1758) 6 22   Sicillaria paestana (Philippi, 1836) 3, 5, 6, 9 12   PUNCTIDAE Puncturu pzymaeum (Draparnaud, 1801) 6 12   Discus rotundabus (O.F. Müller, 1774) 6, 9 12   PRISTILOMATIDAE Vitrea bo	VALLONIIDAE	<i>Acanthinula aculeata</i> (O.F. Muller, 17/4)	8,9	15
CHONDRINDAEGranaria apeniuna (Kuster, 1850)4, 8Chondrina avenacea (Bruguière, 1792)6, 9VERTIGINIDAETruncatellina callicratis (Scacchi, 1833)9IAJaminia quadridens (O.F. Müller, 1774) (small morphotype)7, 8S. 18Jaminia quadridens (O.F. Müller, 1774) (small morphotype)4, 5, 7, 8, 12Merdigera obscura (O.F. Müller, 1774)9SUBULINIDAECecilioides acicula (O.F. Müller, 1774)9SUBULINIDAERumina decollata (Linnacus, 1758)1, 3, 9CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina laminata (Montagu, 1803)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 9Macrogastra plicatula (Draparnaud, 1801)6Clausilia cruciata Studer, 18206PUNCTIDAEPunctum pygmacum (Draparnaud, 1801)6DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 9PRISTILOMATIDAEFirea botterii (Pfeiffer, 1853)4, 6, 8OXYCHILIDAEDaudebardia brevipes (Draparnaud, 1805)6OXYCHILIDAEDaudebardia brevipes (Draparnaud, 1805)6UTTRINDAESemilianceal baoellii (Ta		Gittenbergia sororcula (Benoit, 1857)	6	15
Chondrina avenacea (Bruguere, 1/92) 6, 9   VERTIGINIDAE Truncatellina callicratis (Scaechi, 1833) 9 14   ENDAE Jaminia quadridens (O.F. Müller, 1774) (large morphotype) 7, 8 5, 18   Jaminia quadridens (O.F. Müller, 1774) 6, 8    FERUSSACIIDAE Cecilioides acicula (O.F. Müller, 1774) 9    SUBULINIDAE Rumina decollata (Linnaeus, 1758) 1, 3, 9    CLAUSILIIDAE Medra sp. 6 20   Leucostigma candidescens (Rossmässler, 1835) 1-6, 8, 9, 11, 12 3, 21   Cochlodina laminata (Montagu, 1803) 6 23   Cochlodina laminata (Comparnaud, 1801) 6 23   Cochlodina bidens (Linnaeus, 1758) 6 22   Macrogastra plicathal (Draparnaud, 1801) 6 12   PUNCTIDAE Punctum pygmaeum (Draparnaud, 1801) 6 12   DISCIDAE Discus rotundatus (O.F. Müller, 1774) 6, 8 16   OXYCHILDAE Dundebardia rufg (Draparnaud, 1801) 6 12   DISCIDAE Discus rotundatus (O.F. Müller, 1774) 6, 6	CHONDRINIDAE	Granaria apennina (Kuster, 1850)	4, 8	
VERTIGINIDAE <i>Iruncatellina callicratis</i> (Seacch, 1833)914ENIDAEJaminia quadriders (O.F. Müller, 1774) (small morphotype)7, 85, 18Jaminia quadriders (O.F. Müller, 1774) (large morphotype)4, 5, 7, 8, 124, 19Merdigera obscura (O.F. Müller, 1774)6, 8FERUSSACIIDAECecilioides acicula (O.F. Müller, 1774)9SUBULNIDAERamina decollata (Linnaeus, 1758)11, 3, 9CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1-6, 8, 9, 11, 123, 21Cochlodina bidens (Linnaeus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 9Macrogastra plicatula (Draparnaud, 1801)6Macrogastra plicatula (Draparnaud, 1801)612DISCIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotunadans (O.F. Müller, 1774)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1801)66Mitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia soverbyi (Ferusse, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)6Limax sp. A (black)3, 6, 89Limax sp. A (black)3, 6, 89Limax sp. A (black)3, 6, 89Limax sp. A (black)914HYGROMIIDAEMonacha ef. cantiana (Montagu, 1803)9HYGROMIIDA		Chondrina avenacea (Bruguière, 1792)	6, 9	
ENIDAEJaminia quadridens (O.F. Müller, 1774) (small morphotype)7, 85, 18Jaminia quadridens (O.F. Müller, 1774) (arge morphotype)4, 5, 7, 8, 124, 19Merdigera obscura (O.F. Müller, 1774)6, 8FERUSSACIIDAECecilioides acicula (O.F. Müller, 1774)9SUBULINIDAERumina decollata (Linnaeus, 1758)1, 3, 9CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina bidens (Linnaeus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 92Macrogastra plicatula (Draparnaud, 1801)62PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 91PIISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 8166Oxychilus cf. draparnaud, 1805)666Daudebardia truja (Draparnaud, 1805)6612UTRINIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAELimax sp. A (black)3, 6, 89Limax sp. A (black)3, 6, 891Limax sp. B (brown)611Monacha cf. cantana (Montagu, 1803)91Monacha cf. cantana (Montagu, 1803)91<	VERTIGINIDAE	Truncatellina callicratis (Scacchi, 1833)	9	14
Jamina quadridens (O.F. Müller, 1774) (large morphotype)4, 5, 7, 8, 124, 19Merdigera obscura (O.F. Müller, 1774)6, 8FERUSSACIIDAECecilioides acicula (O.F. Müller, 1774)9SUBULINIDAERumina decollata (Linnacus, 1758)1, 3, 9CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina biens (Linnacus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 99Macrogastra plicatula (Draparnaud, 1801)612Discus rotundatus (O.F. Müller, 1774)6, 912PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 912PRISTILOMATIDAEVitrea buterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia trufa (Draparnaud, 1805)6Daudebardia trufa (Draparnaud, 1805)66Daudebardia soverbyi (Ferusse, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)6LIMAX Sp. A (black)3, 6, 89Limax sp. A (black)3, 6, 89Limax sp. B (brown)66Adora of. cantana (Montagu, 1803)9HYGROMIIDAEMonacha ef. cantana (Montagu, 1803)9HYGROMIIDAEMonacha ef. cantania (Montagu, 1803)9 <trr>Limax sp.</trr>	ENIDAE	Jaminia quadridens (O.F. Müller, 1774) (small morphotype)	7, 8	5, 18
Merdigera obscura (O.F. Müller, 1774)6, 8FERUSSACIIDAECecilioides acicula (O.F. Müller, 1774)9SUBULINIDAERumina decollata (Linnacus, 1758)1, 3, 9CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina bidens (Linnacus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 93Macrogastra plicatula (Draparnaud, 1801)612Olausilia cruciata Studer, 1820612PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 916PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Oxychilus cf. draparnaud, 1805)6612OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66MILACIDAEIandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAELimax sp. A (black)3, 6, 107UTRINIDAELimax sp. A (black)3, 6, 108, 9Limax sp. B (brown)611, 3, 5, 8, 11, 12Cernuellopsis ghisotiti Manganelli et Giusti, 197381HYGROMIIDAECampiaca cf. campiana (Roustia, 1837)101HYGROMIIDAECampiaca f. campianica (Paulucci, 1881)1, 3, 5, 8, 11, 121Cernuellopsis ghisotiti Manganelli e		Jaminia quadridens (O.F. Müller, 1774) (large morphotype)	4, 5, 7, 8, 12	4, 19
FERUSSACIIDAECecilioides acicula (O.F. Müller, 1774)9SUBULINIDAERumina decollata (Linnaeus, 1758)1, 3, 9CLAUSILIIDAEMedora sp.6Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 12Siciliaria paestana (Philippi, 1803)623Cochlodina laminata (Montagu, 1803)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 91Macrogastra plicatula (Draparnaud, 1801)61UNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 91PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 8166OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)666Dudebardia brevipes (Draparnaud, 1805)661UTRINIDAETandonia sowerbyi (Ferussac, 1823)6, 107VITRINIDAELinax cf. maximus Linnaeus, 17583, 6, 108, 9Linax sp. A (black)3, 6, 8911Linax sp. B (brown)6111Cernuella cisalpina (Rosmässler, 1837)1011Cernuella cisalpina (Rosmässler, 1837)1011HYGROMIIDAEMonacha ef. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 121Cernuella cisalpina (Rosmässler, 1837)1011HELICIDAECampralae alf		Merdigera obscura (O.F. Müller, 1774)	6, 8	
SUBULINIDAERumina decollata (Linnaeus, 1758)1, 3, 9CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina bidens (Linnaeus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 99Macrogastra plicatula (Draparnaud, 1801)612DINCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 99PRISTILOMATIDAEVitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)666MILACIDAEZandonia soverbyi (Ferussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66LIMACIDAELimax sp. A (black)3, 6, 89Limax sp. B (brown)644AGRIOLIMACIDAEDeroceras cf. Iolhari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)91Monacha cf. cantiana (Rossmässler, 1837)101HYGROMIIDAECernuellopsis ghistotti Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuellopsis ghistotti Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 121Cernuellopsis ghistotti Manganelli et Giusti, 19881, 4, 5, 7,	FERUSSACIIDAE	Cecilioides acicula (O.F. Müller, 1774)	9	
CLAUSILIIDAEMedora sp.620Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina laminata (Montagu, 1803)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 92Macrogastra plicatula (Draparnaud, 1801)62Clausilia cruciata Studer, 1820612DISCIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 916VITER subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)666Daudebardia rufa (Draparnaud, 1805)666Oxychilus cf. draparnaud, 1805)6612MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)612LIMAX DAELimax ef. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 8911MOnacha ef. cantiana (Montagu, 1803)9111HYGROMIIDAEMonacha ef. cantiana (Montagu, 1803)911HYGROMIIDAEMonacha ef. cantiana (Montagu, 1803)911HELICIDAECampylaea planospira Lamarck, 1822611HELICIDAECampylaea planospira Lamarck, 1822611	SUBULINIDAE	Rumina decollata (Linnaeus, 1758)	1, 3, 9	
Leucostigma candidescens (Rossmässler, 1835)1–6, 8, 9, 11, 123, 21Cochlodina laminata (Montagu, 1803)623Cochlodina bininata (Montagu, 1803)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 922Macrogastra plicatula (Draparnaud, 1801)622Clausilia cruciata Studer, 1820622PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 92PRISTILOMATIDAEFitrea botterii (Pfeiffer, 1853)4, 6, 817Virea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66MILACIDAETandonia soverbyi (Férusae, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66Limax sp. A (black)3, 6, 891Limax sp. A (black)3, 6, 891MIGOMIDAEDeroceras cf. Iothari Giusti, 197381HYGROMIIDAEMonacha cf. camiana (Ponulucci, 1881)1, 3, 5, 8, 11, 121Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 121HELICIDAECampylaea planospira Lamarck, 182261Harrorana signata (Férussac, 1821)6, 9, 1111HELICIDAECampylaea planospira Lamarck, 182261Harrorana signata (Férussac, 1821)6, 9, 1111HE	CLAUSILIIDAE	Medora sp.	6	20
Cochlodina laminata (Montagu, 1803)623Cochlodina bidens (Linnacus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 9Macrogastra plicatula (Draparnaud, 1801)6Clausilia cruciata Studer, 18206PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)6DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 9PRISTILOMATIDAEVitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66OXYCHILIDAEFandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)61UIMACIDAELimax ef. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 891Limax sp. B (brown)666AGRIOLIMACIDAEDeroceras ef. lothari Giusti, 197389HYGROMIIDAEMonacha ef. cantiana (Montagu, 1803)91HYGROMIIDAECernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)101HELICIDAECampylaea planospira Lamarek, 18226Marmorana signata (Férussac, 1821)6, 9, 111HELICIDAECampylaea planospira Lamarek, 18226Marmorana signata (Férussac, 1821)6, 9, 11HELICIDAECamp		Leucostigma candidescens (Rossmässler, 1835)	1-6, 8, 9, 11, 12	3, 21
Cochlodina bidens (Linnaeus, 1758)622Siciliaria paestana (Philippi, 1836)3, 5, 6, 9		Cochlodina laminata (Montagu, 1803)	6	23
Siciliaria paestana (Philippi, 1836)3, 5, 6, 9Macrogastra plicatula (Draparnaud, 1801)6Clausilia cruciata Studer, 18206PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)6DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 9PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817OXYCHILIDAEDaudebardia rufa (Draparnaud, 1801)66OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Oxychilus cf. draparnaud, 1805)66MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66LIMACIDAELimax sp. A (black)3, 6, 89Limax sp. B (brown)666AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 197381HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)911Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 1211Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 1211HYGROMIIDAECamparnada, 1801)111HELICIDAECampanica planospira Lamarck, 1822661Marmorana signata (Férussac, 1821)6, 9, 1111HELICIDAECampalae planospira Lamarck, 1822661Marmorana signata (Férussac, 1821)6, 9, 1111HELICIDAEMarmorana signata (Férussac, 1821)6, 9, 111		Cochlodina bidens (Linnaeus, 1758)	6	22
Macrogastra plicatula (Draparnaud, 1801)6Clausilia cruciata Studer, 18206PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)6DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 9PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)67MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)68LIMACIDAELimax sp. A (black)3, 6, 89Limax sp. A (black)3, 6, 891AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 197381HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)91Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 121Cernuellosis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 121Linac Campanica (Rossmässler, 1837)1011HELICIDAECampylaea planospira Lamarck, 182261Higromia cinctella (Draparnaud, 1801)111HELICIDAECampylaea planospira Lamarck, 182261Higromia cinctella (Draparnaud, 1801)111HELICIDAECantareus apertus (Born, 17741, 3, 12		Siciliaria paestana (Philippi, 1836)	3, 5, 6, 9	
Clausilia cruciata Studer, 18206PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 912PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66Oxychilus cf. draparnaudi (Beck, 1837)2, 6, 8, 107VITRINDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66Limax sp. A (black)3, 6, 899Limax sp. B (brown)666AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 197381HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)91Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 122Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 121HELICIDAECampylaea planospira Lamarck, 182266Marmorana signata (Férussac, 1821)6, 9, 1111HELICIDAECantareus apertus (Born, 17741, 3, 1224		Macrogastra plicatula (Draparnaud, 1801)	6	
PUNCTIDAEPunctum pygmaeum (Draparnaud, 1801)612DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 911PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 8117Vitrea subrimata (Reinhardt, 1871)6, 81616OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)6612MILACIDAEDaudebardia brevipes (Draparnaud, 1805)612MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)612LIMACIDAELimax ef. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 8912Limax sp. B (brown)61212AGRIOLIMACIDAEDeroceras ef. lothari Giusti, 1973814HYGROMIIDAEMonacha ef. cantana (Montagu, 1803)912Monach a ef. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 1212Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 1212LinaxCernuella cisalpina (Rossmässler, 1837)1011HELICIDAECampylaea planospira Lamarck, 1822614Marmorana signata (Férussac, 1821)6, 9, 1113, 1224		Clausilia cruciata Studer, 1820	6	
DISCIDAEDiscus rotundatus (O.F. Müller, 1774)6, 9PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66Oxychilus cf. draparnaudi (Beck, 1837)2, 6, 8, 107MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66LIMACIDAELimax ef. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 896AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 197386HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)99Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 122Cernuella cisalpina (Rossmässler, 1837)101HELICIDAECampylaea planospira Lamarck, 182266Marmorana signata (Férussac, 1821)6, 9, 116Helix liegato OF, Müller, 17741, 3, 1224	PUNCTIDAE	Punctum pygmaeum (Draparnaud, 1801)	6	12
PRISTILOMATIDAEVitrea botterii (Pfeiffer, 1853)4, 6, 817Vitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66LIMACIDAELimax ef. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 896AGRIOLIMACIDAEDeroceras ef. lothari Giusti, 197386HYGROMIIDAEMonacha ef. cantiana (Montagu, 1803)99Monacha ef. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 121Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 121HELICIDAECampanica (Rossmässler, 1837)101HELICIDAECampylaea planospira Lamarck, 182262Helix ligata O.F. Müller, 17741, 3, 1224	DISCIDAE	Discus rotundatus (O.F. Müller, 1774)	6, 9	
Vitrea subrimata (Reinhardt, 1871)6, 816OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)66Daudebardia brevipes (Draparnaud, 1805)66Oxychilus cf. draparnaudi (Beck, 1837)2, 6, 8,107MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)66LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 896AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 197387HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)99Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 121Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 121HELICIDAECampylaea planospira Lamarck, 182266Marmorana signata (Férussac, 1821)6, 9, 117Helix ligata O.F. Müller, 17741, 3, 1224	PRISTILOMATIDAE	Vitrea botterii (Pfeiffer, 1853)	4, 6, 8	17
OXYCHILIDAEDaudebardia rufa (Draparnaud, 1805)6Daudebardia brevipes (Draparnaud, 1805)6Oxychilus cf. draparnaudi (Beck, 1837)2, 6, 8, 10MILACIDAETandonia sowerbyi (Férussac, 1823)6, 10VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)6LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 89Limax sp. B (brown)66AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12HLICIDAECernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Helix ligata O.F. Müller, 17741, 3, 1224		Vitrea subrimata (Reinhardt, 1871)	6, 8	16
Daudebardia brevipes (Draparnaud, 1805)6Oxychilus cf. draparnaudi (Beck, 1837)2, 6, 8,10MILACIDAETandonia sowerbyi (Férussac, 1823)6, 10VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)6LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 89Limax sp. B (brown)66AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O,F, Müller, 17741, 3, 12	OXYCHILIDAE	Daudebardia rufa (Draparnaud, 1805)	6	6
Oxychilus cf. draparnaudi (Beck, 1837)2, 6, 8,10MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)61LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 899Limax sp. B (brown)66AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 12		Daudebardia brevipes (Draparnaud, 1805)	6	
MILACIDAETandonia sowerbyi (Férussac, 1823)6, 107VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)6LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 89Limax sp. B (brown)6AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)924		Oxychilus cf. draparnaudi (Beck, 1837)	2, 6, 8,10	
VITRINIDAESemilimacella bonellii (Targioni Tozzetti, 1873)6LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 89Limax sp. B (brown)6AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 12	MILACIDAE	Tandonia sowerbyi (Férussac, 1823)	6, 10	7
LIMACIDAELimax cf. maximus Linnaeus, 17583, 6, 108, 9Limax sp. A (black)3, 6, 89Limax sp. B (brown)6AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 12	VITRINIDAE	Semilimacella bonellii (Targioni Tozzetti, 1873)	6	
Limax sp. A (black)3, 6, 89Limax sp. B (brown)6AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224	LIMACIDAE	Limax cf. maximus Linnaeus, 1758	3, 6, 10	8, 9
Limax sp. B (brown)6AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224		Limax sp. A (black)	3, 6, 8	9
AGRIOLIMACIDAEDeroceras cf. lothari Giusti, 19738HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10HELICIDAECampylaea planospira Lamarck, 1822Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224		<i>Limax</i> sp. B (brown)	6	
HYGROMIIDAEMonacha cf. cantiana (Montagu, 1803)9Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10Hygromia cinctella (Draparnaud, 1801)1HELICIDAECampylaea planospira Lamarck, 1822Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224	AGRIOLIMACIDAE	Deroceras cf. lothari Giusti, 1973	8	
Monacha cf. campanica (Paulucci, 1881)1, 3, 5, 8, 11, 12Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10Hygromia cinctella (Draparnaud, 1801)1HELICIDAECampylaea planospira Lamarck, 1822Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224	HYGROMIIDAE	Monacha cf. cantiana (Montagu, 1803)	9	
Cernuellopsis ghisottii Manganelli et Giusti, 19881, 4, 5, 7, 8, 11, 12Cernuella cisalpina (Rossmässler, 1837)10Hygromia cinctella (Draparnaud, 1801)1HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224		Monacha cf. campanica (Paulucci, 1881)	1, 3, 5, 8, 11, 12	
Cernuella cisalpina (Rossmässler, 1837)10Hygromia cinctella (Draparnaud, 1801)1HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224		Cernuellopsis ghisottii Manganelli et Giusti, 1988	1, 4, 5, 7, 8, 11, 12	
Hygromia cinctella (Draparnaud, 1801)1HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224		Cernuella cisalpina (Rossmässler, 1837)	10	
HELICIDAECampylaea planospira Lamarck, 18226Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 1224		Hygromia cinctella (Draparnaud, 1801)	1	
Marmorana signata (Férussac, 1821)6, 9, 11Cantareus apertus (Born, 1778)9Helix ligata O.F. Müller, 17741, 3, 12	HELICIDAE	<i>Campylaea planospira</i> Lamarck. 1822	6	
Cantareus apertus (Born, 1778) 9   Helix ligata O.F. Müller, 1774 1, 3, 12 24		Marmorana signata (Férussac. 1821)	6, 9, 11	
Helix ligata O.F. Müller. 1774 1. 3. 12 24		Cantareus apertus (Born, 1778)	9	
		Helix ligata O.F. Müller, 1774	1. 3. 12	24

Table 2. List of the species recorded in the sampled stations, Lepini Mountains (southern Latium, Italy).



Figures 2–9. Terrestrial gastropods from Lepini Mountains. Fig. 2: *Cochlostoma* cf. *adamii*. Fig. 3: *Leucostigma candidescens*. Fig. 4: *Jaminia quadridens* (large morphotype). Fig. 5: *Jaminia quadridens* (small morphotype). Fig. 6: *Daudebardia rufa*. Fig. 7: *Tandonia sowerbyi*. Fig. 8: *Limax* cf. *maximus*. Fig. 9: *Limax* sp. A (black) and L. cf. *maximus*.



Figures 10–17. Terrestrial gastropods from Lepini Mountains. Fig. 10: *Platyla similis*. Fig. 11: *Acicula* sp. Fig. 12: *Punctum pygmaeum*. Fig. 13: *Acanthinula aculeata*. Fig. 14: *Truncatellina callicratis*. Fig. 15: *Gittenbergia sororcula*. Fig. 16: *Vitrea subrimata*. Fig. 17: *Vitrea botterii*. Scale bar: 1 mm.



Figures 18–23. Terrestrial gastropods from Lepini Mountains. Fig. 18: *Jaminia quadridens* (small morphotype). Fig. 19: *Jaminia quadridens* (large morphotype). Fig. 20: *Medora* sp. Fig. 21: *Leucostigma candidescens*. Fig. 22: *Cochlodina bidens*. Fig. 23: *Cochlodina laminata*. Scale bar: 5 mm (whole specimens); 2.5 mm (details).

black specimens of *Limax* Linnaeus, 1758 was found in different stations, also in sympatry with *L*. cf. *maximus* (Fig. 9). A further *Limax* species with uniform brown colour was recorded.

A small black specimen of *Deroceras* Rafinesque, 1820 was collected in Stn. 8, externally resembling *Deroceras lothari*, a species described as endemic to Reatini Mountains (northen Latium) (Giusti, 1973). It is provisionally reported as *Deroceras* cf. *lothari* pending further study.

Holm oak and mixed woods are poorer in terms of both species and number of specimens. Pastures show a relatively reduced number of species as well, some of which deserve some comments. *Jaminia quadridens* occurs in two distinct forms, here referred to as small and large morphotype, respectively (Figs. 4, 5, 19, 20). At a conchological level, they mainly differ with regard to shell size. Measurements on over 60 specimens show that these two forms can be readily distinguished by the shell width, which shows a clear gap between the two morphotypes (Fig. 25). Moreover, the large morphotype seems to show a proportionally larger width. However, this feature needs confirmation due to the small amount of recorded specimens of the small morphotype. The large morphotype is widespread and was recorded from several stations. Conversely, the small morphotype is uncommon and was recorded only at higher altitudes. A similar scenario with different morphs of *J. quadridens* occurring syntopically have been recorded also in other localities of Central Apennines, often with the occurrence of an additional, dextral morph.

*Monacha campanica*, which we regard as a distinct species, is found throughout the Liri Valley up to high altitude. The genitalia of *M. campanica* are characterized by a very short penis provided with a large penial papilla and a slender epiphallus. The flagellum is as long as the epiphallus. The bursa copulatrix shows a thick duct, widened at the base. Specimens from Lepini Mountains share the same anatomy but show a rather different shell. In fact, they are smaller (max. diam. 16 mm) and lighter than those described by the marquise Paulucci (1881) (max. diam. 21 mm). Moreover, they show a less depressed spire and a narrower umbilicus. As a consequence, they are reported in Table 2 as *Monacha* cf. *campanica*, pending further study.

In Stn. 9 (Bassiano, road to Semprevisa, near the spring) an empty shell clearly different from



Figure 24. Helix ligata, height 35 mm, with genitalia (scale bar: 5 mm).



Fig. 25. Height and width measurements of *Jaminia quadridens* s.l. specimens from Lepini Mountains. Best-fit lines for the two morphotypes are shown.

*Monaca* cf. *campanica* has been collected. It is provisionally reported as *Monacha* cf. *cantiana* waiting for anatomical data.

*Cernuellopsis ghisottii* is the most abundant species in grasslands and pastures above 1000 m. This species shows a disjunct distribution in the Apennines. Southern populations occur in the Pollino (Calabria-Basilicata) and Sirino (Basilicata) massifs and extend up to Alburni Mountains (Campania), wherease populations from Central Apennines are frequent on the western coastal chains and extend up to Simbruini Mountains (Latium). The species was never recorded from intermediate mountains of central-northen Campania.

*Helix ligata* is widespread in Lepini Mountains but never abundant. Different populations show a variable appearance. Specimens from Carpinetana Valley were genetically studied by Fiorentino et al. (2016) and are very similar to specimens from Apennine valley floors showing a yellowish background due to the presence of periostracum, whereas specimens from Stn. 1 (pass to Campo di Segni) living in exposed pasture with stones show a whitish background (Fig. 24) likely due to the loss of periostracum that make it resemble *Helix delpretiana* Paulucci, 1878 (Giusti, 1973). However, the anatomy of specimens from Stn. 1 corresponds to *Helix ligata*, even though they are genetically distinct from populations from Central Apennines (Fiorentino et al., 2016).

### CONCLUSIONS

The preliminary checklist here presented shows that at least 43 species of terrestrial gastropods occur in the Lepini Mountains. The richest environment is represented by beech forests, with 28 recorded species. Along with species already reported from Central Apennines, a new isolated population of Medora is recorded. Due to the extremely limited distribution, this population can be considered vulnerable and likely in need of protection. Jaminia quadridens occurs in two clearly distinct morphs without intermediate forms, mainly differing in their size. Further research is required in order to ascertain whether they actually belong to different species. Some species remain undetermined, whereas others were determined by comparison, pending further research.

## REFERENCES

- Amori G., Corsetti L. & Esposito C, 2002. Mammiferi dei Monti Lepini. Quaderni di Conservazione della Natura, 11, Ministero dell'Ambiente e della Tutela del Territorio - Istituto Nazionale per la Fauna Selvatica "Alessandro Ghigi".
- Bodon M. & Cianfanelli S., 2008. Una nuova specie di *Platyla* per il sud Italia (Gastropoda: Prosobranchia: Aciculidae). Bollettino Malacologico, 44: 27–37.
- Bodon M., Favilli L., Giannuzzi Savelli R., Giovine F., Giusti F., Manganelli G., Melone G., Oliverio M., Sabelli B. & Spada G., 1995. Gastropoda Prosobranchia, Heterobranchia Heterostropha. In: Minelli A., Ruffo S. & La Posta S. (Eds), Checklist delle specie della fauna italiana 14, Calderini, Bologna, 60 pp.
- Bouchet Ph. & Rocroi J.P., 2005. Classification and nomenclator of gastropod families. Malacologia, 47: 1–397.
- Colomba M.S., Liberto F., Reitano A., Renda W., Pocaterra G., Gregorini A. & Sparacio I., 2012. Molecular studies on the genus *Medora* H. et A. Adams, 1855 from Italy (Gastropoda Pulmonata Clausiliidae). Biodiversity Journal, 3: 571–582.
- Corsetti L., Angelini C., Copiz R., Mattoccia M. & Nardi G., 2015. Biodiversità dei Monti Lepini. Edizioni Belvedere, Latina, 352 pp.

- Fehér Z., Deli T. & Sólymos P., 2010. Revision of *Granaria frumentum* (Draparnaud 1801) (Mollusca, Gastropoda, Chondrinidae) subspecies occurring in the eastern part of the species' range. Journal of Conchology, 40: 201–217.
- Fiorentino V., Manganelli G., Giusti F. & Ketmaier V., 2016. Recent expansion and relic survival: Phylogeography of the land snail genus *Helix* (Mollusca, Gastropoda) from south to north Europe. Molecular Phylogenetics and Evolution, 98: 358–372.
- Giusti F., 1973. Notulae malacologicae XVI. I molluschi terrestri e di acqua dolce viventi sul massiccio dei Monti Reatini (Appennino Centrale). Lavori della Società Italiana di Biogeografia, NS 2: 423–574.
- Giusti F., Castagnolo L. & Manganelli G., 1985. La fauna malacologica delle faggete italiane: brevi cenni di ecologia, elenco delle specie e chiavi di riconoscimento dei generi e delle entità più comuni. Bollettino Malacologico, 21: 69–144.
- Giusti F., Grappelli C., Manganelli G., Fondi R. & Bulli L., 1986. An attempt of natural classification of the genus *Medora* in Italy and Yugoslavia, on the basis of conchological, anatomical and allozymic characters (Pulmonata: Clausiliidae). Lavori della Società Italiana di Malacologia, 22: 259–341.
- Hallgass A. & Vannozzi A., 2014. The continental molluscs from Mount Circeo (Latium, Italy). Biodiversity Journal, 5: 151–164.
- ICZN, 2015. Opinion 2355 (Case 3581): Turbo bidens Linnaeus, 1758 (Gastropoda, Clausillidae): request to set aside the neotype not granted. Bulletin of Zoological Nomenclature, 72: 159–161.
- Kadolsky D., 2009. *Turbo bidens* Linnaeus 1758 (Gastropoda: Clausiliidae) misidentified for 250 years. Journal of Conchology, 40: 19–30.
- Latella L., 1995. La fauna cavernicola dei Monti Lepini. Notiziario del Circolo Speleologico Romano, NS 6-7, 1991-'92: 78–119.
- Magrini P., 2005. Un nuovo *Neobacanius* anoftalmo del Lazio (Insecta, Coleoptera: Histeridae). Aldrovandia, 1: 55–62.
- Manganelli G., Bodon M., Favilli L. & Giusti F., 1995. Gastropoda Pulmonata. In: Minelli A., Ruffo S. & La

Posta S. (Eds.), Checklist delle specie della fauna italiana 16, Bologna (Calderini): 1–60.

- Manganelli G., Bodon M., Favilli L., Castagnolo L. & Giusti F., 1998. Checklist delle specie della fauna italiana, molluschi terrestri e d'acqua dolce. Errata e addenda. Bollettino Malacologico, 33: 151–156.
- Manganelli G., Bodon M. & Giusti F., 2000. Checklist delle specie della fauna italiana, molluschi terrestri e d'acqua dolce. Errata e addenda. Bollettino Malacologico, 36: 125–130.
- Nardi G., Di Russo C. & Latella L., 2002. Populations of *Nepa cinerea* (Heteroptera: Nepidae) from hypogean sulfurous water in the Lepini Mountains (Latium, Central Italy). Entomological News, 113: 125–130.
- Nordsieck H., 2012. Ergänzung der Revision der Gattung Medora H. & A. Adams: Die Medora-Arten Italiens (Gastropoda, Stylommatophora, Clausiliidae, Alopiinae), mit Beschreibung einer neuen Unterart von Medora dalmatina Rossmässler. Conchylia, 42: 75– 81.
- Pace R., 1975. An exceptional endogeous beetle: Crowsoniella relicta n. gen. n. sp. of Archostemata Tetraphaleridae from central Italy. Bollettino del Museo Civico di Storia Naturale di Verona, 2: 445– 458.
- Paulucci M., 1881. Contribuzione alla fauna malacologica italiana. Specie raccolte dal Dott. C. Cavanna negli anni 1878, 1879, 1880 con elenco delle conchiglie abruzzesi e descrizione di due nuove *Succinea*. Bullettino della Società Malacogica Italiana, 7: 69–180.
- Sani F., Del Ventisette C., Montanari D., Coli M., Nafissi P. & Piazzini A., 2004. Tectonic evolution of the internal sector of the Central Apennines, Italy. Marine and Petroleum Geology, 21: 1235–1254.
- Sbordoni V., 1971. Il popolamento animale e vegetale dell'Appennino Centrale. Lavori della Società Italiana di Biogeografia, NS 2: 595–614.
- Zallot E., Groenenberg D.S.J., De Mattia W., Fehér Z. & Gittenberger E., 2015. Genera, subgenera and species of the Cochlostomatidae (Gastropoda, Caenogastropoda, Cochlostomatidae). Basteria 78: 63–88.