

## A Taxonomic study of the Pygmygrasshopper (Orthoptera Tetrigidae) from India with description of a new species

Sunil Kumar Gupta & Kailash Chandra

Zoological Survey of India, Prani Vigyan Bhawan, 'M' Block New Alipore, Kolkata-700053, West Bengal, India; e-mail: skumarento@gmail.com

### ABSTRACT

A new species of genus *Coptotettix* Bolivar, 1887, *C. magedhensis* n. sp. (Orthoptera Tetrigidae), is described from Jagdalpur, Chhattisgarh, India. Also studied with 19 species belonging to 7 genera under two subfamilies Scelimeninae and Tetriginae under family Tetrigidae.

### KEY WORDS

Orthoptera; Tetriginae; new species; Scelimeninae; Barnawapara; Durg; Kabirdham.

Received 03.05.2017; accepted 11.06.2017; printed 30.06.2017

### INTRODUCTION

Scattered information about the occurrence of Orthoptera fauna from India had been reported by several scientists: Kirby (1914) described 5 new genera and 10 new species, besides the already known 23 genera and 73 species. Hancock (1912, 1915) studied the Indian Tetrigidae. Fletcher (1921) gave a full list of synonyms and references along with the distribution for the Tetrigidae of India and the adjacent countries. Hebard (1929) studied the South Indian Tetrigids. Günther (1935-1941) revised the subfamilies of Tetrigidae with the exception of Batrachidinae and Tetriginae. Steinmann (1964-1971a, b) compiled the taxonomical work of the family Tetrigidae in the form of Checklist to give a zoogeographical picture of the species of the world. Grant (1966) studied the genera of Batrachideinae. Shishodia (1991a) studied 'Taxonomy and zoogeography of the Tetrigidae of North eastern India' and 77 species were dealt with, of which 13 species were new, 6 were new records for India, one new genus was proposed and two new synonyms were established. Shishodia (1984, 1987 a,b, 1991a,b,c, 1993, 1995, 2000a, 2007a), Paranjape &

Bhalerao (1986), Paranjape et al. (1987), Wagan & Kevan (1992) mainly worked on the taxonomy of Tetrigoidea in India.

A little work has been done about pygmy grasshopper fauna from Chhattisgarh state i.e. Shishodia (2000b) reported 77 species of crickets and grasshoppers from Bastar, of which 13 species of Tetrigidae family. Recently Skejo & Gupta (2015) gave a specific status of *Hedotettix cristatus* Karny, 1915. Gupta (2016 a, b) worked on Orthoptera fauna of Chhattisgarh and described two species of Family Tetrigidae: *Euparatettix dandakaranyensis* Gupta, 2016, *Ergatettix subtruncatus* Gupta, 2016; Gupta et al. (2016) described also *Hedotettix angulatus* Gupta, Shi et Chandra, 2016.

The genus *Coptotettix* Bolivar, 1887 is the most diverse genus of Tetrigidae erected by Bolivar (1887). This genus includes 79 species and subspecies (Eades et al., 2017), of which, 12 species reported from India (Shishodia et al., 2010).

During the extensive and intensive survey of these areas from 2011 to 2014, a total of 20 species belonging to 7 genera under 2 subfamilies Scelimeninae and Tetriginae under family Tetrigidae with a new species were collected from different

localities from Chhattisgarh and the results are shown below.

## MATERIAL AND METHODS

The survey was carried out in Chhattisgarh state, covering four districts Durg, Jagdalpur, Korba and Surguja and seven Protected areas i.e. Sitanadi Wildlife Sanctuary, Udanti Wildlife Sanctuary, Achanakmar Wildlife Sanctuary, Gurughasidas National Park, Badalkhol Wildlife Sanctuary, Barnawapara Wildlife Sanctuary and Bhoramdev Wildlife Sanctuary.

The specimens were collected by sweeping over pebbles on the banks of rivers, by using an insect net and killed by benzene vapor in killing jar. The specimens were preserved dry and pinned and they were studied under a Leica stereozoom Microscope (Leica M205 A) and photographs were taken using the software Leica Application Suite (LAS V3.8). All the measurements are given in millimetres. The type specimens were deposited in the Central Entomological Laboratory (CEL) of the Zoological Survey of India in Kolkata while the other specimens in the National Zoological Collection of the Zoological Survey of India, Kolkata.

All the specimens were collected by S.K. Gupta and colleagues.

ABBREVIATIONS. DC: Day collection; FRH: Forest Rest House; NC: Night collection; NP: National Park; WLS: Wildlife Sanctuary.

## RESULTS

### *Systematics*

Order ORTHOPTERA Olivier, 1789  
 Suborder CAELIFERA Ander, 1939  
 Superfamily TETRIGOIDEA Ramber, 1838  
 Family TETRIGIDAE Ramber, 1838  
 Subfamily SCeliminenae Ramber, 1838  
 Genus *Thoradonta* Hancock, 1909

#### 1. *Thoradonta apiculata* Hancock, 1915

*Thoradonta apiculata* - Hancock, 1915: 8  
*Thoradonta apiculata* - Shishodia, 2000a: 233

EXAMINED MATERIAL. Chhattisgarh, Dhamtari, Sitanadi WLS, Sankra FRH, 01.VII.2012, 1 female, NC; Durg, Badbhumi FRH, 28.III.2014, 1 male, NC; Bilaspur, Achanakmar WLS, Chhaparwa FRH, 10.VI. 2012, 1 female, NC.

DISTRIBUTION. India: Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Chhattisgarh, Manipur, Meghalaya, Orissa, Tamil Nadu and West Bengal. Elsewhere: Myanmar.

#### 2. *Thoradonta spiculoba* Hancock, 1912

*Thoradonta spiculoba* - Hancock, 1912: 138  
*Thoradonta pruthii* - Shishodia, 2000b: 27-80.

EXAMINED MATERIAL. Chhattisgarh, Korba, Telimnala, 03.VI.2012, 1 male, DC, Jashpur, Badalkhol WLS; Bachhram beat, 25.V.2012, 9 females, DC; Dhamtari, Sitanadi WLS; Gidhava, 21.X.2012, 1 female, DC; Koriya, Gurughasidas NP, Sonhut FRH, 19.III.2014, 1 male, NC; Durg, Badbhumi FRH, 28.III.2014, 1 male, DC; Surguja, Taravayaya Nala, 17.IX.2012, 1 female.

DISTRIBUTION. India: Chhattisgarh, Karnataka, Madhya Pradesh, Maharashtra, Orissa and West Bengal.

#### 3. *Thoradonta nodulosa* (Stål, 1861)

*Tetrix nodulosa* - Stål, 1861: 346  
*Thoradonta nodulosa* - Shishodia, 2000a: 233

EXAMINED MATERIAL. Chhattisgarh, Jashpur, Badalkhol WLS; Bachhram beat, 25.V.2012, 1 male, DC; Koriya, Gurughasidas NP, Sonhut FRH, 19.III.2014, 1 male, NC.

DISTRIBUTION. India: Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Himachal Pradesh, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Sikkim, Tamil Nadu, Tripura and West Bengal. Elsewhere : Borneo, China, Hainan, Java, Malaysia, Myanmar, Perak, Pinang, Singapore, Sri Lanka, Sumba and Sumatra.

#### 4. *Thoradonta bengalensis* Shishodia, 1991

*Thoradonta bengalensis* - Shishodia, 1991a: 34  
*Thoradonta bengalensis* - Shishodia, 2003: 82.

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Gabod, 08.X.2011, 1 male, DC; Parbatiya Nala, 04.VI.2012, 1 male, NC; Domar pani, 24.IV.2012, 1 female, DC; Jagdalpur, Sonarpal beat, 17.X.2011, 3 males, DC; Dhamtari, Sitanadi WLS; Sitanadi river, 1 female, DC, 17.X.2011, 1 female, DC; Kabirdham, Bhoramdev WLS; Rajadhar Forest, 01.XII.2011, 1 female, DC; Jashpur, Badalkhol WLS; Bachhram beat, 25.V.2012, 11 females, DC; Korba, Kudmura nala, 31.V.2012, 1 male, DC; Bilaspur, Achanakmar WLS, Karidongri FRH, 19.IX.2012, 1 female, DC; Gariyaband, Udanti WLS; Kulhadighat, 28.VI.2014, 1 male, DC.

DISTRIBUTION. India: Chhattisgarh and West Bengal.

Genus *Euscelimena* Günther, 1938

#### 5. *Euscelimena harpago* (Serville, 1838)

*Tetrix harpago* - Serville, 1838: 763  
*Scelimena uncinata* - Shishodia, 1991b: 34  
*Euscelimena harpago* - Kulkarni & Shishodia, 2005: 326.

EXAMINED MATERIAL. Chhattiagarh, Raigarh, Gomerda WLS; Kanakabira, 19.II.2012, 2 males, 3 females, DC; Gariyaband, Udanti WLS; Jhariyahara, 24.VI.2014, 3 males, 5 females, DC; 29.VI.2014, 1 female, DC.

DISTRIBUTION. India: Andhra Pradesh, Bihar, Karnataka, Gujarat, Madhya Pradesh, Maharashtra, Orissa, Tamil Nadu and Uttar Pradesh. Elsewhere: Sri Lanka.

Subfamily TETRIGINAE Serville, 1838  
 Genus *Paratettix* Bolivar, 1887

#### 6. *Paratettix histrionicus* (Stål, 1861)

*Tetrix histrionicus* - Stål, 1861: 347  
*Euparatettix histrionicus* - Shishodia, 1987b: 93  
*Paratettix histrionicus* - Storozhenko & Dawwrueng, 2015: 530.

EXAMINED MATERIAL. Durg, Durg camp, 23.VIII.2011, 1 male, DC; Rampur, 24.VIII.2011,

6 females, DC; Raipur, Barnawapara WLS; Domarpani, 28.VI.2012, 1 female, DC; Nawapara, 19.VII.2013, 1 female, DC; Dhamtari, Sitanadi WLS; Gidhava, 21.X.2012, 2 females, DC; Gahnasiyar, 20.VIII.2013, 1 male, DC.

DISTRIBUTION. India: Andaman and Nicobar Islands, Andhra Pradesh, Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Delhi, Himachal Pradesh, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Orissa, Rajasthan, Tamil Nadu, Tripura and West Bengal. Elsewhere: Australia, Borneo, Caledonia, Celebes, East Africa, East Afghanistan, Holland, Indonesia, Iran, Java, Malaysia, Myanmar, Nepal, New Ireland, Pakistan, Philippines, Queensland, Saudi Arabia, Solomon Islands, South China, Sumatra, Sri Lanka and Taiwan.

#### 7. *Paratettix cingalensis* (Walker, 1871)

*Tettix cingalensis* - Walker, 1871: 827  
*Paratettix variegatus* - Bolivar, 1887: 280  
*Paratettix cingalensis* - Chandra et al., 2007: 2686

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Barnawapara, 04.IX.2011, 1 female, NC; 06.VII.2011, 1 male, NC.

DISTRIBUTION. India: Arunachal Pradesh, Assam, Chhattisgarh, Goa, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Meghalaya, Manipur, Orissa, Sikkim, Tamil Nadu and Tripura. Elsewhere: Borneo, Hainan, Malaysia, Java, Philippines, Sri Lanka, Sumatra and Taiwan.

Genus *Euparatettix* Hancock, 1904

#### 8. *Euparatettix nodulosus* Hancock, 1912

*Euparatettix nodulosus* - Hancock, 1912: 154  
*Ergatettix nodulosus* - Hebard, 1929: 587  
*Ergatettix guentheri* - Steinmann, 1970a: 234  
*Euparatettix nodulosus* - Gupta, 2016a: 121

EXAMINED MATERIAL. Chhattisgarh, Koriya Gurughasidas NP; 28.IX.2011, 1 male, DC; Uravpara, 24.X.2011, 1 male, DC; Raipur, Barnawapara WLS; Gabod, 08.X.2011, 1 male, DC; Latadar, 01.VII.2013, 1 male, DC; Durg, Badbhum FRH, 28.III.2014, 1 male, 1 female, NC; Saldevi mandir, 25.IX.2012, 1 male, DC.

DISTRIBUTION. India: Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Himachal Pradesh, Madhya Pradesh, Maharashtra, Meghalaya, Mizoram, Orissa, Tripura, Uttar Pradesh and West Bengal. Elsewhere: Bangladesh, Nepal and Sri Lanka.

Genus *Ergatettix* Kirby, 1914

**9. *Ergatettix callosus* (Hancock, 1915)**

*Indatettix callosus* - Hancock, 1915: 130

*Indatettix callosus* - Fletcher, 1921: 35

*Ergatettix callosus* - Shishodia, 1993: 205

EXAMINED MATERIAL. Chhattisgarh, Jagdalpur, Bhanpuri Forest, 20.X.2011, 1 male, DC; Ulnar village, 16.III.2012, 1 male, DC; Raipur, Barnawapara WLS; Hardi Nala, 29.VI.2013, 1 male, DC; Lalbandh Pond, 13.VII.2013, 1 male, DC.

DISTRIBUTION. India: Chhattisgarh and West Bengal.

**10. *Ergatettix interruptus* (Brunner von Wattenwyl, 1893)**

*Paratettix interruptus* - Brunner von Wattenwyl, 1893: 104

*Euparatettix interruptus* - Kirby, 1910: 31

*Indatettix interruptus* - Hancock, 1915: 130

*Ergatettix interruptus* - Blackith, 1992: 54

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Bar forest, 29.V.2013, 1 male, DC; Dhamtari, Sitanadi WLS; Bhainsa sankra, 30.VIII.2013, 1 male, DC.

DISTRIBUTION. India: Chhattisgarh. Elsewhere: Laos, Myanmar and Thailand.

**11. *Ergatettix dorsifera* (Walker, 1871)**

*Tettix dorsifera* - Walker, 1871: 825

*Ergatettix dorsiferus* - Shishodia, 2000: 21

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Gabod, 08.X.2011, 1 male, DC; Hardi Pathar, 29.III.2012, 1 male, DC; 30.III.2012, 1 male, DC; Keduva Nala, 03.VI.2013, 1 male, DC; Hardi Nala, 26.VI.2013, 1 male, DC;

Latadadar, 28.VI.2013, 1 male, DC; Durg; Rampur, 24.VIII.2011, 1 female, DC; Jagdalpur, Magrapara, 06.I.2012, 1 female, DC; Belguda, 18.I.2012, 1 female, DC; Malgaon, 09.III.2012, 1 female, DC; Surguja, Tara Rest House, 16.IX.2012, 1 male, DC; Koriya; Gurughasidas NP; Uravpara, 24.X.2011, 1 male, DC.

DISTRIBUTION. India: Andaman & Nicobar Islands, Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Delhi, Himachal Pradesh, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Orissa, Rajasthan, Sikkim, Tripura, Tamil Nadu, Uttarakhand, Uttar Pradesh and West Bengal. Elsewhere: Afghanistan, Bangladesh, China, Central Asia, Java, Greater Sunda Islands, Iran, Indonesia, Myanmar, Nepal, Sri Lanka, Sumatra, Taiwan and Turkmenistan.

Genus *Coptotettix* Bolivar, 1887

**12. *Coptotettix tricarinatus* Shishodia, 1991**

*Coptotettix tricarinatus* - Shishodia, 1991a: 169

*Coptotettix tricarinatus* - Shishodia, 2003: 83


EXAMINED MATERIAL. Chhattisgarh, Jagdalpur, Malgaon, 09.III.2012, 1 male, DC; Koriya Gurughasidas NP; Sonhut FRH, 05.V.2013, 1 male, NC.

DISTRIBUTION. India: Chhattisgarh and West Bengal.

**13. *Coptotettix magedhensis* n. sp. (Figs. 1–4)**

EXAMINED MATERIAL. Holotype, female, DC. India, Chhattisgarh, Jagdalpur, Makdi, Magedha, 09.XI.2011, 19°05'45.6"N, 82°03'9.3"E, 592 m, legit S.K. Gupta and colleagues, Registration no. 18713/H5. Paratypes: 1 female, same data as holotype. Registration no. 187714/H5. 1 female, Jagdalpur, Hathguda, 29.III.2012, 190545.6N, 82039.3E, 561 m. Registration no. 187715/H5.

DESCRIPTION OF HOLOTYPE. Female. Vertex narrower than a compound eye, little elevated, median carina of the vertex present in the distal half of the length of vertex, elevated, fossulae present. Lateral carinae of the vertex present, strongly elevated, with tubercles. Median ocellus situated below the level of the lower margin of the compound eyes, in the place where facial carina ends. Lateral ocelli situ-



Figures 1–4. *Coptotettix magedhensis* n. sp., holotype female. Fig. 1: dorsal view.  
Fig. 2: lateral view. Fig. 3: Paranota and Tegmen. Fig. 4: Ovipositor.

ated in the level of the distal margin of the compound eyes. Frontal costa bifurcation starts in about upper  $\frac{1}{4}$  of the compound eye height and runs parallel to the median ocelli forming a wider scutellum. Antennal grooves are situated at the level of the lower margin of compound eyes or slightly below it. Eyes in frontal view sub-globular, in lateral view sub-globular. Eyes can be not close to the pronotum so the occipital area is visible.

Pronotum. Pronotum granulated with fine, medium sized and large granules, wrinkled. Anterior margin truncate, extended upto the last abdominal segment, acute at apex. Median carina is continuous towards the whole length of pronotum bearing 6–7 small projections of variable height and size. External carinae fused to humero-apical carinae. Humero-apical carinae forming with external lateral carinae obtuse angle, two lateral lobes present. Posterior margin of lateral lobe truncated.

Wings. Tegmina elongated, wings present, visible under pronotum extend up to 4th or 5th abdominal segments.

Fore legs: femora 3 times as long as wide, Upper and lower margins of femora straight; serration on external face indistinct; tibia clothed with 7 spines and some small hairs; tibial serration on external face distinct, second tarsal segments 5 times longer than first.

Mid legs: femora 3.6 times as long as wide, serrated carina on external face distinct, Upper and lower margins of femora slightly undulate, upper and lower mid carinae with several black warts, tibiae clothed with 4 spines and some hairs; second tarsal segments 2.9 times longer than first.

Hind legs: Hind femora robust 2.6 times as long as wide, upper margins serrated, tibiae with 8 outer and 5 inner spines. first tarsal segments 1.5 times longer than third; tip of pulvilli I and II spinulately produced and pulvilli III obtusely, pulvilli II distinctly longer than length of I and also II pulvilli. Subgenital plate in ventral view square shaped; Ovipositor elongated. Epiproct as long as wide near the base, with obtuse apex. Cerci cylindrical, wide near the base with obtuse tip.

Coloration. Body color from brown to dark blackish. Pronotal tubercles usually dark black. Fore tibiae, mid tibiae and hind tibiae with two dark brown hues, genicular lobe of hind femora dark brown; fore legs of III tarsal segment with dark

black brown hue. Ventral region of hind femora dark black color. Tip of spines dark blackish.

Measurements (all in millimeters). Body length Female. (from the tip of the vertex to the end of the abdomen): 8.959; head length: 0.944; Vertex: 0.582. compound eye length: 0.659, width: 0.391. Antenna length: 2.994. Antenna groove length: 0.145, width: 0.223, scapes length: 0.219, pedicel length: 0.177, scutellum width: 0.212. Pronotum length: 7.691, shoulder width: 2.383. Fore leg: femur length: 1.535, width: 0.501; tibia length: 1.760, width: 0.214; tarsus length (by segments): I: 0.114, II: 0.577; claw length: 0.192. Mid leg: femur length 1.776, width 0.482; tibial length 1.692, width 0.262; tarsus length: I: 0.184, II: 0.538; claw length 0.287. Hind leg: femur length: 4.856, width: 1.807; tibia length: 3.808, width: 0.236; tarsus length: I: 0.712, II: 0.132, III: 0.465; pulvilli length: I: 0.245, II: 0.226, III: 0.312; claw length 0.226. Abdomen length: 4.886. Wings: tegmen length: 1.393, width: 0.450; Dorsal ovipositor valves length: 1.252, height: 0.302; ventral ovipositor valves length: 1.048, height: 0.149; cerci length: 0.310, width: 0.125.

VARIABILITY. The paratypes female have no substantial morphological differences from the holotype described. Male unknown.

ETYMOLOGY. The specific epithet refers to the type locality.

DISTRIBUTION. *Coptotettix magedhensis* n. sp. is known actually from the type locality only: India, Chhattisgarh.

Genus *Criotettix* Bolivar, 1887

#### 14. *Criotettix bispinosus* (Dalman, 1818)

*Acrydium bispinosum* - Dalman, 1818: 77

*Criotettix bispinosus* - Günther, 1937: 123

*Criotettix bispinosus* - Shishodia, 2000a: 232

EXAMINED MATERIAL. Chhattisgarh, Koriya, Gurughasidas NP, Ghatoriya Pahad, 28.XI.2013, 1 male, DC; Thitima Pahar, 24.X.2013, 1 male, 1 female, DC.

DISTRIBUTION. India: Arunachal Pradesh, Assam, Bihar, Chhattisgarh, Madhya Pradesh, Manipur, Meghalaya, Tripura and West Bengal. Else-

where: Borneo, Celebes, China, Hainan, Hong kong, Java, Luzon, Malaysia, Myanmar, Sulawesi, Sumatra, Taiwan, Thailand and Vietnam.

Genus *Hedotettix* Bolívar, 1887

### 15. *Hedotettix attenuatus* Hancock, 1904

*Hedotettix attenuatus* - Hancock, 1904: 151

*Hedotettix attenuatus* - Kirby, 1914: 73

*Hedotettix attenuatus* - Chandra et al., 2007: 2687

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Barnawapara, 24.VII.2011, 1 male, 1 female, NC; Raigarh, Gomerda WLS; Naresh Nagar, 24.VIII.2011, 1 male, NC; Surguja, Anjani Beat, 10.XI.2011, 1 female, DC; Korba, Paharjamri Beat, 24.XI.2011, 1 female, DC; Bijakhara, 01.VI.2012, 1 male, DC; Koriya, Gurughasidas NP; Sonhut Forest Rest House, 21.IV.2013, 1 male, 2 females, NC; 05.V.2013, 1 female, NC; 19.III.2014, 1 female, NC; Kamrajee, Bagridanar, 22.IX.2013, 1 male, DC; Thitima Pahar, 24.X.2013, 1 male, DC; Berdev Nala, 17.XI.2013, 1 male, DC; Ghatoriya Pahar, 28.XI.2013, 1 male, DC; Rewala, 09.XII.2013, 1 male, 2 females, DC; Durg, Badbhum FRH, 08.II.2014, 1 female, NC; 31.III.2014, 1 female, NC; Gariyaband, Udanti WLS; Ganga Jamuna village, 28.VI.2014, 1 male, 2 females, NC.

DISTRIBUTION. India: Andaman and Nicobar Islands, Arunachal Pradesh, Assam, Chhattisgarh, Himachal Pradesh, Manipur, Meghalaya, Orissa, Sikkim, Tripura, Uttarakhand, Uttar Pradesh and West Bengal. Elsewhere: Bangladesh, Celebes, China, Java, Myanmar, Pakistan, Sri Lanka, Sulawesi, Sumatra, Taiwan, Thailand and Vietnam.

### 16. *Hedotettix gracilis* (Haan, 1842)

*Acridium gracile* - Haan, 1842: 169

*Hedotettix gracilis* - Hancock, 1915: 123.

*Hedotettix gracilis* - Chandra et al., 2007: 2687

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Barnawapara, 10.VII.2011, 1 male, DC; 14.VII.2011, 2 females, NC; 20.VII.2011, 1 female, NC; 24.VII.2011, 1 female, NC; 09.XI.2011, 1 male, DC; 03.II.2012, 1 female, NC; Keduva Nala, 09.VII.2011, 1 female, DC; Dulilibahal, Forest, 10.IX.2011, 1 male, DC; Latadar,

02.I.2012, 1 female, DC; Surguja, Bengehi Pahar, 10.XI.2011, 1 male, DC; Kabirdham, Bhoramdev WLS; Sitalpani, 19.XI.2011, 1 male, DC; Sankri River, 30.IX.2013, 1 male, DC; Raigarh, Gomerda WLS; Gomerda village, 28.V.2013, 1 male, DC; Koriya, Gurughasidas NP; Rawanpat, 23.X.2013, 1 male, DC.

DISTRIBUTION. India: Andaman and Nicobar Islands, Arunachal Pradesh, Assam, Chhattisgarh, Himachal Pradesh, Karnataka, Kerala, Madhya Pradesh, Maharashtra, Manipur, Meghalaya, Orissa, Rajasthan, Sikkim, Tripura, Uttarakhand, Uttar Pradesh and West Bengal. Elsewhere: Bangladesh, Celebes, China, Java, Myanmar, Pakistan, Sri Lanka, Sulawesi, Sumatra, Taiwan, Thailand and Vietnam.

### 17. *Hedotettix costatus* Hancock, 1912

*Hedotettix costatus* - Hancock, 1912: 147

*Hedotettix costatus* - Shishodia & Tandon, 2000: 221

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Latadar, 27.IX.2011, 1 female, NC.

DISTRIBUTION. India: Assam, Bihar, Himachal Pradesh, Meghalaya, Sikkim, Tripura, Uttar Pradesh and West Bengal. Elsewhere: Bangladesh, Nepal and Sulawesi.

### 18. *Hedotettix punctatus* Hancock, 1909

*Hedotettix punctatus* - Hancock, 1909: 422

*Hedotettix punctatus* - Shishodia, 2007: 89

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Gabod, 08.X.2011, 1 male, DC; Koriya, Gurughasidas NP; Dev Nala, 09.III.2013, 1 male, DC; Durg, Badbhum FRH, 30.III.2014, 1 male, DC; Kabirdham, Bhoramdev WLS; Sankri River, 30.IX.2013, 2 females, DC; Gariyaband, Udanti WLS; Jhariyahabara, 22.II.2014, 1 male, DC.

DISTRIBUTION. India: Chhattisgarh, Karnataka, Madhya Pradesh, Maharashtra and Rajasthan. Elsewhere: Borneo and Himalayan region.

### 19. *Hedotettix grossus* Hancock, 1915

*Hedotettix grossus* - Hancock, 1915: 124

*Hedotettix grossus* - Shishodia, 2000: 243

EXAMINED MATERIAL. Chhattisgarh, Raipur, Barnawapara WLS; Nawapara, 23.VII.2011, 1 female, NC; Kabirdham, Bhoramdev WLS; Kherpani Forest, 16.XI.2011, 1 female, NC; Sankri river, 30.IX.2013, 1 female, NC; Dhamtari, Sitanadi WLS; Bhainsa Sankra, 04.IX.2013, 1 female, DC; Sankra FRH, 24.II.2014, 1 female, NC.

DISTRIBUTION. India: Arunachal Pradesh, Assam, Chhattisgarh, Manipur, Meghalaya, Tripura and West Bengal.

### 20. *Hedotettix quadriplagiatus* (Walker, 1871)

*Tettix quadriplagiatus* - Walker, 1871: 828

*Hedotettix quadriplagiatus* - Blackith, 1992: 88

EXAMINED MATERIAL. Chhattisgarh, Kabirdham, Bhoramdev WLS; Sitalpani, 19.XI.2011, 1 male, DC; Rajadhar Forest, 01.XII.2011, 1 female, DC; Gariyaband, Udanti WLS; Jhariyabahara, 24.IV.2014, 1 male, 3 females, DC; Durg, Badbhum FRH, 31.III.2014, 2 females, DC.

DISTRIBUTION. India: Chhattisgarh and North India.

### ACKNOWLEDGEMENTS

The authors are thankful to Director, Zoological Survey of India, Kolkata for providing necessary facilities and encouragement. The authors are thankful to CAMPA (Compensatory Afforestation Fund Management and Planning Authority) for the funding. The first author is also grateful to Dr Suresh Chand, Officer-in-charge, Orthoptera section and all staff Orthoptera section for their support.

### REFERENCES

- Bolívar I., 1887. Essai sur les acridiens de la trib Tettigidae. Annales de la Société entomologique belge, 31: 175–313.
- Blackith R.E., 1992. Tetrinidae (Insecta; Orthoptera) of Southeast Asia: Annotated catalogue with partial translated keys and bibliography. 1–248, key LI–LIV.

- Brunner V.W., 1893. Revision du systeme de Orthopteres et descriptions des especes rapportees par M. Leonardo Fea de Birmanie. Annali del Museo Civico di Storia Naturale "Giacomo Doria" di Genova, 2, 13(33): 1–230.
- Chandra K., Gupta S.K. & Shishodia M.S., 2007. A Check-list of Orthoptera of Madhya Pradesh and Chhattisgarh. Zoos' Print Journal, 22: 2683–2687.
- Dalman J.W., 1818. Nagra nya Genera och Species af Insekter beskrifna. Kongliga Svenska Vetenskaps-Akademien Handlingar, 39, 69–89.
- Eades D.C., Otte D., Cigliano M.M. & Braun H., 2017. Orthoptera Species File. Version 5.0/5.0. [accessed on 3rd may 2017] <http://Orthoptera.SpeciesFile.org>
- Fletcher T.B., 1921. Catalogue of Indian Insects. pt. I Acrydiidae (Tettigidae) pp. iv+40.
- Grant H.J. Jr., 1966. The pacific genera of the subfamily Batrachideinae (Orthoptera : Tetrigidae). Pacific Insects, 8: 579–601.
- Günther K., 1935. Acrydiinae von den key Inseln (Orthoptera: Acrididae). Eos, 11: 109–115.
- Günther K., 1937. Acrydiinae (Orthoptera: Acrididae) von Java der Kleinen Sunda Inselns und Nordaustralien. Revue suisse de zoologie, 44: 121–143.
- Günther K., 1941. Revision der Acrydiinae ausbeute H. Sauters von Formosa (Orth.). Stettin Entomological Society, 102: 145–165.
- Gupta S.K., 2016a. *Euparatettix dandakaranyensis* sp. nov. (Tetrigidae: Tetriginae) - a new pygmy grasshopper species from Central India. Zootaxa, 4097: 118–124. <http://dx.doi.org/10.11646/zootaxa.4097.1>
- Gupta S.K., 2016b. A new pygmy grasshopper species (Tetrigidae: Tetriginae) from Central India. Zootaxa. 4097: 280–286. <http://dx.doi.org/10.11646/zootaxa.4097.2>
- Gupta S.K., Shi J.P & Chandra K., 2016. *Hedotettix angulatus* sp. nov. (Orthoptera: Tetrigidae: Tetriginae) a new pigmy grasshopper from, Central India. Zootaxa, 4173: 466–475. <http://doi.org/10.11646/zootaxa.4173.5.3>
- Haan W. De, 1842. Bijdragen tot de Kentnis der Orthoptera. In: Temminck C.J., Verh. Natuur 1. Geschiedenis van de Nederlandse Overzee. Bezitt, 2: 45–232.
- Hancock J.L., 1904. The Tettigidae of Ceylon. Spolia zeylanica, 2: 97–157.
- Hancock J.L., 1909. Further studies of the Tetriginae (Orthoptera) in the Oxford University Museum. Transactions of the Entomological Society of London, 387–426.
- Hancock J.L., 1912. Tetriginae (Acridiinae) in the Agricultural Research Institute, Pusa, Bihar, with description of new species. Memoirs of the Department of Agriculture in India. Entomological series, 4: 131–160.

- Hancock J.L., 1915. Indian Tetriginae (Acridiinae). Records of the Indian Museum, 11: 55–135.
- Hebard M., 1929. Acridiinae (Orthoptera, Acrididae) of Southern India. Revue suisse de zoologie, 36: 565–592.
- Kirby W.F., 1914. The fauna of British India, including Ceylon and Burma, Orthoptera (Acridiidae), London, 1: IX + 276 pp.
- Kirby W.F., 1910. A Synonymic Catalogue of Orthoptera. 3. Orthoptera Saltatoria part 2, Locustidae vel Acrididae, 1–299.
- Kulkarni P.P. & Shishodia M.S., 2005. Insecta : Orthoptera. Conservation Area Series, 24: Fauna of Melghat Tiger Reserve, Zoological Survey of India: 317–340.
- Paranjape S.Y. & Bhalerao A.M., 1986. Biosystematic and ethoecological studies on family Tetrigidae (Orthoptera). Proceedings of the 3rd Oriental Entomology Symposium. February 21–24, 1984. Volume 1. Association for the Advancement of Entomology, University of Kerala. Kariavattom, India: 19–23.
- Paranjape S.Y., Bhalerao A.M. & Naidu N.M., 1987. On etho-ecological characteristics and phylogeny of Tetrigidae. In: Baccetti B. (Ed.), Evolutionary biology of orthopteroid insects. Ellis Horwood Ltd, Chichester: 386–395.
- Serville J.G.A., 1838. Histoire naturelle des insectes. Orthoptères. I–XVIII, 1–776, 14 plates. DOI: <http://dx.doi.org/10.5962/bhl.title.95609>
- Shishodia M.S., 1984. A new genus *Neocoptotettix* from India (Orthoptera: Tetrigidae). Oriental Insurance, 18: 13–16.
- Shishodia M.S., 1987a. A new combination proposed for Indian Grouse-Locusts (Orthoptera: Tetrigidae). Bulletin Zoological Survey of India, 8: 131–134.
- Shishodia M.S., 1987b. Tetrigidae: Orthoptera : Insecta (Grouse-locusts). State Fauna Series, 1: Fauna of Orissa, Part 1. Zoological Survey of India: 91–102.
- Shishodia M.S., 1991a. Taxonomy and Zoogeography of the Tetrigidae (Orthoptera : Tetridoidea) of North Eastern India. Records of the Zoological Survey of India, Occasional Paper, 140: vi + 1–203.
- Shishodia M.S., 1991b. On a collection of Grouse-Locusts (Orthoptera: Tetrigidae) from Maharashtra, India, with some new distributional records. Records of the Zoological Survey of India, 88: 29–36.
- Shishodia M.S., 1991c. Grouse Locusts (Insecta: Orthoptera: Tetrigidae) of the Kanha National Park, District-Mandla, Madhya Pradesh, India. Records of the Zoological Survey of India, 89: 101–104.
- Shishodia M.S., 1993. Insecta: Orthoptera : Tetridoidea. State Fauna Series, 3: Fauna of West Bengal, Part 4. Zoological Survey of India: 179–226.
- Shishodia M.S., 1995. Insecta: Orthoptera: Tetridoidea. State Fauna Series, 4: Fauna of Meghalaya, Part 3. Zoological Survey of India: 165–208.
- Shishodia M.S., 2000a. Insecta: Orthoptera : Tetridoidea. State Fauna Series, 7: Fauna of Tripura, Part 2. Zoological Survey of India: 231–246.
- Shishodia M.S., 2000b. Short and long horned grasshoppers and crickets of Bastar district, M.P., India. Records of the Zoological Survey of India, 98: 27–80.
- Shishodia M.S., 2003. Genera and species of Orthopteran insects. Bionotes, 5: 82–84.
- Shishodia M.S., 2007. Insecta: Orthoptera: Tetrigidae. State Fauna Series, 14: Fauna of Mizoram. Zoological Survey of India: 217–224.
- Shishodia M.S. & Tandon S.K., 2000. Insecta: Orthoptera: Acridoidea and Eumastacoidea. State Fauna Series, 7: Fauna of Tripura, Part 2. Zoological Survey of India: 197–230.
- Shishodia M.S., Chandra K. & Gupta S.K., 2010. An annotated checklist of Orthoptera (Insecta) from India. Records of the Zoological Survey of India, Occasional Paper, 314, 1–366.
- Skejo J. & Gupta S.K., 2015. On the specific status of *Hedotettix cristatus* Karny, 1915 (Tetrigidae: Tetriginae). Zootaxa. 4018: 584–592. DOI: <http://dx.doi.org/10.11646/zootaxa.4018.4.7>
- Stål C., 1861. Orthoptera species novas descripsit. Kongliga Svenska Fregatten Eugenies Resa Omkring Jorden, 3: 299–350. Stockholm [1860]
- Steinmann H., 1964. Some new Tetrigid species and subspecies from Asia (Orthoptera: Tetrigidae). Acta zoologica Academiae Scientiarum Hungaricae, 10: 457–468.
- Steinmann H., 1969. The Tetrigidae (Orthoptera) of the Neogaea. Folia entomologica hungarica, 22: 383–404.
- Steinmann H., 1970a. Check-list of the Tetricidae (Orthoptera) of the Oriental Faunal region. Acta zoologica Academiae Scientiarum Hungaricae, 16: 215–240.
- Steinmann H., 1970b. The Tetricidae (Orthoptera) of the Notogaea. Opuscula Zoologica (Budapest), 10: 155–164.
- Steinmann H., 1971a. The Tetricids of the Nearctic subregion. Acta zoologica Academiae Scientiarum Hungaricae, 17: 381–385.
- Steinmann H., 1971b. The Tetricidae (Orthoptera) of the Palaeartic fauna. Folia entomologica hungarica, 24: 323–332.
- Storozhenko S.YU. & Dawwrueng P., 2015. New and little-known pygmy grasshoppers (Orthoptera: Tetrigidae) from Thailand. Zootaxa, 4052, 527–554. DOI: <http://dx.doi.org/10.11646/zootaxa.4052.5.2>
- Wagan M.S. & Kevan D.K. McE., 1992. Studies on some

Tetrigidae (Orthoptera) from India, Pakistan and Sri Lanka. *Tropical Zoology*, 5: 167–194.

Walker F., 1871. Supplement to the Catalogue of Dermaptera Saltatoria. London, 116 pp.