28

www.biodiversityjournal.com

ISSN 2039-0394 (Print Edition) ISSN 2039-0408 (Online Edition)

with the support of



Biodiversity Journal

SEPTEMBER 2017, 8 (3): 773-868

FOR NATURALISTIC RESEARCH AND ENVIRONMENTAL STUDIES



Eupteryx petasitidis Ferrari, 1882 (Auchenorrhyncha Cicadellidae). The taxon from Piedmont has been described as Eupteryx carpini petasitidis. Later, it was considered a synonym of E. atropunctata (Goeze, 1778). Recently (2014), E. petasitidis was re-established as a valid species based on a detailed study of coloration and genital morphology. Until now, this species has been found only in the Italian mainland (from Piedmont to Calabria).

Apparently, it is monophagous and lives on *Petasites hybridus* (L.) Gaertn., Mey. and Scherb., and has been found only in rather low altitudes (100 - 800 m) on their host plant in shadowy places near rivers or brooks. Its closest relatives are *E. atropunctata* (Goeze, 1778) and *E. aurata* (Linnaeus, 1758), which both are polyphagous with a wide range of host plants and present in many European countries.

Christoph Bückle. Neckarhalde 48, D-72070 Tübingen, Germany; e-mail: rosaurusrex@gmx.de (photos by G. Kunz)