

Two new records of fossil gastropods from the Italian upper Pleistocene

Mauro M. Brunetti

Calle Navas 106, 14511 Navas del Selpillar, Spain; e-mail: mbrunetti45@gmail.com

ABSTRACT

The examination of numerous materials related to the upper Pleistocene (“Tirreniano” sensu Authors) from Sardinia and Calabria led to the discovery of two new fossil species for the Mediterranean basin: *Sinum bifasciatum* (Récluz, 1851) (Gastropoda Naticidae), found both in the upper Pleistocene of Sardinia and in that of Calabria, and *Morula nodulosa* (C.B. Adams, 1845) (Gastropoda Muricidae), found in the Sardinian upper Pleistocene. Both species are discussed and illustrated in this paper.

KEY WORDS

Sinum; *Morula*; Pleistocene; Tyrrhenian.

Received 10.09.2018; accepted 29.10.2018; printed 30.12.2018; published online 07.01.2019

INTRODUCTION

The study of material coming from the upper Pleistocene of the Mediterranean basin led to the discovery of two new fossil gastropod molluscs for the Mediterranean basin: *Sinum bifasciatum* (Récluz, 1851) and *Morula nodulosa* (C.B. Adams, 1845). Both species come from deposits with the typical hot fauna with Senegalese affinity to *Persistrombus latus* Gmelin, 1791 (= *Strombus bubonius* Lamarck, 1822), *Monoplex trigonus* (Gmelin, 1791) [= *Cymathium ficoides* (Reeve, 1844)], *Gemophos viveratus* (Kiener, 1834), *Conus ermineus* Born, 1778, (= *Conus testudinarius* Hwass in Bruguière, 1792), etc. (Gignoux, 1913; Trevisan & Di Napoli, 1938; Mirigliano, 1953; Segre, 1954; Malatesta, 1954; Ruggieri & Buccheri, 1968; Settepassi, 1971; Ruggieri & Melone, 1975; Vazzana, 2008).

MATERIAL AND METHODS

The samples, collected during surface searches, come from different deposits of the upper Pleis-

tocene from Sardinia and Calabria (see Examined material). The taxonomy follows Worms (2018) and the others cited publications.

ABBREVIATIONS. H: maximum height of the shell, measured from the apex up to the front end of the siphonic channel; L: maximum width of the shell; ex: specimen/s; CMF: Maurizio Forli collection; CPC: Paolo Crovato collection; CMB: Mauro M. Brunetti collection.

RESULTS

Systematics

Classis GASTROPODA Cuvier, 1795
Subclassis CAENOGASTROPODA Cox, 1960
Ordo LITTORINIMORPHA Golikov et Starobogatov, 1975

Superfamilia NATICOIDEA Guilding, 1834
Familia NATICIDAE Guilding, 1834
Genus *Sinum* Röding, 1798
Type species: *Sinum haliotoideum* Linnaeus, 1758

Sinum bifasciatum (Récluz, 1851) (Figs. 1–3)

Sigaretus bifasciatus Récluz, 1851: 190, tab. 6, figs. 3–4

Sigaretus philippii Weinkauff, 1883: 23, tab. 5, figs. 1–3

Sinum bifasciatum Settepassi, 1972, tab. 2

Sinum bifasciatum Ardonini & Cossignani, 2004: 119

Sinum bifasciatum Rolán, 2005: 91, fig. 383

Sinum bifasciatum Giannuzzi Savelli et al., 1997: 202, figs. 831a-f.

EXAMINED MATERIAL. Porto Alabe (Nuoro, Italy) “Tirreniano”, 1 ex, CMB; Trumbacà (Reggio Calabria, Italy) “Tirreniano”, 1 ex, CMF; Saracinello (Reggio Calabria, Italy) “Tirreniano”, 4 ex, CPC.

DESCRIPTION. Shell oval, depressed, not very robust. Maximum width of the shells (L): 17.5 mm (Porto Alabe) and 11.2 mm (Trumbacà). Protoconch composed of three whorls. Teleoconch largely formed by the last lap, very wide. Sculpture composed of numerous spiral cords separated by narrow furrows. Umbilicus absent.

DISTRIBUTION. *Sinum bifasciatum* has a very wide geographical distribution, ranging from the western Mediterranean Sea (Giannuzzi-Savelli et al., 2002) to all of West Africa, up to Angola (Ardovini & Cossignani, 2004; Rolán, 2005).

REMARKS. For *S. bifasciatum* it is the first record as a fossil for the Mediterranean basin. Among the two currently known species of the genus *Sinum* present in the Pliocene of the Mediterranean basin, *S. bifasciatum* bears some resemblance to *S. striatum* (De Serres, 1829) (Fig. 3). From this species, *S. bifasciatum* differs mainly for the different spiral sculpture as specified also by Pedriali & Robba (2009): “*The Mediterranean and West African species Sinum bifasciatum (Récluz, 1851) has a larger, 3 whorled protoconch whith rapidly expanding last whorl, a more depressed teleoconch and narrowed spirals separated by shallow, narrow grooves*”. *Sinum bifasciatum* shows greater differences with the other Pliocene species, *S. perregulare* (Sacco, 1891) (see figures published in Pedriali & Robba, 2009), of smaller dimensions and with the presence of an evident umbilicus.

Ordo NEOGASTROPODA Wenz, 1938

Superfamilia MURICOIDEA Rafinesque, 1815

Familia MURICIDAE Rafinesque, 1815

Genus *Morula* Schumacher, 1817

Type species: *Morula papillosa* Schumacher, 1817

Morula nodulosa (C.B. Adams, 1845) - Fig. 4

Purpura nodulosa C.B. Adams, 1845: 2

Ricinula ferruginea Reeve, 1846: 50, tab. 1–6.

Morula nodulosa Houart, 1997: 83, figs. 174–175, 197, 256–260

Morula nodulosa Ávila et al., 2002: 354, figs. 100–103

Morula nodulosa Ardonini & Cossignani, 2004: 155–156


Morula nodulosa Rolán, 2005: 129, figs. 532–533.

EXAMINED MATERIAL. Porto Alabe (Nuoro) “Tirreniano”, 2 ex, CMB.

DESCRIPTION. Shell sturdy biconoidal, with multispiral protoconch, smooth. Maximum height of the shell (H): 8 mm. Teleoconch with 5 convex whorls, sculpture robust, formed by large roundish nodules originating from the crossing of the strong axial ribs with the spiral cords. The oval opening has 4–5 robust round teeth on the outer lip, 2–3 poorly visible teeth are present on the columellar side. Short siphonal channel, slightly bent to the left. Umbilicus absent.

DISTRIBUTION. This species has a very wide, amphilantic geographical distribution, from the Caribbean to the whole of West Africa to Angola (Houart, 1997; Rolán, 2005). Reported as a fossil of the Upper Pleistocene of the Azores Islands (Garcia-Talavera, 1990; Ávila et al., 2002, 2008).

REMARKS. This is the first record of *M. nodulosa* as a fossil for the Mediterranean basin. Houart (1997) illustrated the protoconch, the radula and the typical material of this species that comes from Jamaica. The reports of *M. nodulosa* as a fossil of the upper Pleistocene are all related to the island of Santa Maria, the Azores archipelago (Garcia-Talavera, 1990; Ávila et al., 2002, 2008). Ávila et al. (2008) report *M. nodulosa* as belonging to a group of species that left the Azores during the last glaciation, ascribing it to the genus *Trachypollia* Woodring, 1928. The material from the Sardinian Pleistocene is smaller than today’s populations.



Figures 1, 2. *Sinum bifasciatum* (Récluz, 1851). Fig. 1: Porto Alabe (Nuoro, Italy) "Tirreniano", L = 17.5 mm, CMB. Fig. 2: Trumbacà (Reggio Calabria) "Tirreniano", detail of the sculpture, L = 11.2 mm, CMF. Figure 3. *Sinum striatum* (De Serres, 1829) Poggio alla Staffa (Siena, Italy), Zanclean, detail of the sculpture, L = 16 mm, CMB. Figure 4. *Morula nodulosa* (C.B. Adams, 1845) Porto Alabe (Nuoro) "Tirreniano", H = 8 mm, CMB.

ACKNOWLEDGEMENTS

We would like to thank Maurizio Forli (Prato, Italy) and Paolo Crovato (Napoli, Italy) for their help in this work.

REFERENCES

- Ardovini R. & Cossignani T., 2004. Conchiglie dell'Africa Occidentale. L'Informatore piceno, Ancona, 320 pp.
- Ávila S.P., Rui A., Azevedo M.N., Cachão M. & Garcia-Talavera F., 2002. Checklist of the Pleistocene marine Molluscs of Prainha and Laghoinas (Santa Maria Island, Azores). Açoreana, 9: 343–370.
- Ávila S.P., Madeira P., Mendes N., Rebelo A., Medeiros A., Gomes C., Garcia-Talavera F., da Silva C.M., Cachão M., Hillaire-Marcel C. & de Frias Martins A.M., 2008. Mass extinctions in the Azores during the last glaciation: fact or myth? Journal of Biogeography, 35: 1123–1129.
- Garcia-Talavera F., 1990. Fauna tropical en el Neotirreniano de Santa Maria (I. Azores). Lavori S.I.M., 23: 439–443.
- Giannuzzi-Savelli R., Pusateri F., Palmeri A. & Ebreo C., 2002. Atlante delle conchiglie marine del Mediterraneo vol. 2. Edizioni Evolver, Roma, 258 pp.
- Gignoux M., 1913. Les formations marines pliocénées et quaternaires de l'Italie du Sud et de la Sicilie. Thèses Faculté des Sciences de l'Université de Lyon, I A. Rey, Lyon, 393 pp.
- Houart R., 1997. Les Muricidae d'Afrique Occidentale. II. Ocenebrinae, Ergalatinae, Tripterotyphinae, Typhinae, Trophoninae & Rapaninae. Apex, 12: 49–91.
- Malatesta A., 1954. Fossili delle spiagge Tirreniane. Bollettino del Servizio Geologico d'Italia, 76: 9–17.
- Mirigliano G., 1953. La macrofauna del Tirreniano di Gallipoli (Lecce). Italian Journal of Zoology, 20: 115–122.
- Pedriali L. & Robba E., 2009. A revision of the Pliocene Naticids of northern and central Italy. III. The Sub-families Poliniceinae and Sininae. Rivista Italiana di Paleontologia e Stratigrafia, 115: 371–429.
- Récluz C.A., 1851. Catalogue des espèces du genre Sigaret (*Sigaretus* Lk). Journal de Conchyliologie, 2: 163–191, pl. 6.
- Reeve L.A., 1846. Monograph of the genus *Ricinula*. In: Conchologia Iconica, vol. 3, pl. 1–6 and unpaginated text. L. Reeve & Co., London.
- Rolán E., 2005. Malacological fauna from the Capo Verde Archipelago. Part 1 Polyplacophora and Gastropoda. Conchbooks, Hackenheim, 455 pp.
- Ruggieri G. & Buccheri G., 1968. Una Malacofauna tirreniana dell'isola di Ustica (Sicilia). Geologica Romana, 7: 27–58.
- Ruggieri G. & Melone G., 1975. La malacofauna del Tirreniano di Tommaso Natale (Palermo). Bollettino Società Paleontologica Italiana, 12: 217–222.
- Segre A.G., 1954. Il Tirreniano del golfo di Terranova Pausania (Olbia) e la sua fauna malacologica. Bollettino del Servizio Geologico d'Italia, 76: 43–84.
- Settepassi F., 1971. Atlante malacologico I molluschi marini viventi nel Mediterraneo. Vol. 2, INIVAG, Roma, 250 pp.
- Settepassi F., 1972. Atlante malacologico I molluschi marini viventi nel Mediterraneo. Vol. 3, INIVAG, 250 pp.
- Trevisan L. & Di Napoli E., 1938. Tirreniano, Siciliano e Calabriano nella Sicilia sudoccidentale. Estratto Giornale di Scienze Naturali ed Economiche, 39: 1–39.
- Vazzana A., 2008. *Ranilia constricta* (Milne Edwards, 1880) nel Tirreniano di Trumbacà vicino Reggio Calabria (Decapoda Brachyura Raninidae). Il Naturalista siciliano, 32: 381–388.
- Weinkauf H.C., 1883. Die Gattung *Sigaretus*. In: Systematisches Conchylien-Cabinet von Martini und Chemnitz, 2nd ed. (Küster H.C., Ed.). 6: 1–50.
- World Register of Marine Species: WoRMS, 2018. <http://www.marinespecies.org/> - Last access: gg.mm.aaaa