

## ***Prodidomus amaranthinus* (Lucas, 1846) (Araneae Gnaphosidae) new for the Italian Araneofauna**

Antonino Dentici<sup>\*1</sup> & Franco Ciro Amata<sup>2</sup>

<sup>1</sup>via E.Cialdini 2, 90124 Palermo, Italy; e-mail: a.dentici@virgilio.it

<sup>2</sup>Piazza S.Antonino 1, 94018 Troina (Enna), Italy; e-mail: amata.francociro@gmail.com

\*Corresponding author

### **ABSTRACT**

The spider *Prodidomus amaranthinus* (Lucas, 1846) (Araneae Gnaphosidae), species never reported before for Italy, was found, sampled and identified in Ribera province of Agrigento, in Sicily (Italy). This is the first record of the genus and species for Sicily and Italy. Additional notes on taxonomy, distribution and biology are provided.

### **KEY WORDS**

Araneae; Spider; Araneofauna; Gnaphosidae; Sicily.

Received 25.03.2021; accepted 16.04.2021; published online 28.04.2021

### **INTRODUCTION**

The genus *Prodidomus* Hentz, 1874 belong to family Gnaphosidae (Prodidominae), transferred from the Prodidomidae to the Gnaphosidae (Prodidominae) by Azevedo et al. (2018).

Of the 54 species belonging to genus *Prodidomus* (WSC, 2021), only 7 are known in the Mediterranean area (Netwing et al., 2021): *P. amaranthinus* (Lucas, 1846), *P. flavidus* (Simon, 1844), *P. geniculatus* Dalmás, 1919, *P. hispanicus* Dalmás, 1919, *P. redikorzevi* Spassky, 1940, *P. rollasoni* Cooke, 1964 and *P. rufus* Hentz, 1847 (although this species, reported for Spain, is identified in “cfr.” (Ferrández & Carrillo, 2018).

Among these, *P. amaranthinus*, reported for Algeria (Lucas, 1846; Walckenaer, 1847; Pickard-Cambridge, 1872; Dalmás, 1919), Egypt (Simon, 1880; Simon, 1907), Libya (Cooke, 1964), Morocco (Simon, 1870; Dalmás, 1919), Tunisia (Dalmás, 1919; Cooke, 1964), and more recently for Cyprus (Bosman et al., 2019), Greece/Crete (Bosman et al., 2013), Portogallo (Pérez & Blasco, 1986; Branco et al., 2019), Spagna (Pérez & Blasco,

1986; Branco et al., 2019) and Turkey (Asia) (Topçu & Türkeş, 2010), is the species with the widest Mediterranean distribution.

Cooke (1964) writes “*is the most widespread species of the family, and occurs along the whole length of the North African coast from Morocco to Syria*”. This distribution contrasts strongly with the isolated occurrence of most other species, and is presumably due to its tolerance of a “*wider range of environments than other members of the family*”.

### **MATERIAL AND METHODS**

On February 2, 2021, the spiders (1 male, 1 female and 2 juveniles) were collected in an area, with low vegetation and sporadic trees, located in Ribera (37°25'18.0" N 13°16'49.2" E), Agrigento, (Sicily, Italy). Three other females specimens were collected on 21 March 2021, in Ribera.

They were photographed with a Canon EOS 7D Mark II with a EF 100 mm Macro with 20 mm extension rings and 1.4 canon multiplier (F.C. Amata).

All the specimens were sampled and preserved in centrifuge tube, fixed in 75% ethanol (Levi, 1977). Observation with a stereomicroscope allowed a correct determination, which follow Pérez & Blasco (1986). They are currently stored in the collection of one of the authors (A. Dentici, Palermo, Italy).

## RESULTS

### Systematics

Ordo ARANEAE Clerck, 1757

Familia GNAPHOSIDAE Pocock. 1898

Genus *Prodidomus* Hentz, 1847



Figures 1, 2. *Prodidomus amaranthinus* from Sicily: male (Fig. 1) and female (Fig. 2).

### *Prodidomus amaranthinus* (Lucas, 1846)

**DESCRIPTION.** The specimens found in Sicily, belonging to group 1 of Dalmas (1919), coincide with the descriptions of Cooke (1964) and Perez & Blasco (1986). In particular, they show a wine-red coloration of the opisthosoma, with sparse hair, and of the opisthosoma, darker in the female specimens, greyish brown thickly covered with hair. The length of the body ranges from 6 to 9 mm, with the females being larger than the males. Peculiar are the ocular distribution, with dark anterior median eyes in contrast with the remaining pearly ones, of which the anterior line slightly curved and the posterior one extremely procured, to create as a whole an almost triangular shape and the spinnerets, of cylindrical shape truncated with those posterior, clearly larger than the anterior, squat, compactly arranged and thickly covered with bristles.

**DISTRIBUTION AND BIOLOGY.** Mediterranean. As reported by Cooke (1964), although the species is associated with desert or very arid environments, it has been observed in very different contexts, such as *Quercus* woods “a habitat which, during the winter months at least, must be very damp”, or for example “At Tabarka, the stone beneath which *P. amaranthinus* was found was lying on damp mud covered with long grass, and being a steep north-facing slope, conditions were such that I doubt whether the site would ever be entirely dry at any time of year”. These observations suggest a strong adaptability of the species to different environmental conditions, which explains the wide distribution of the species.

In Sicily, *P. amaranthinus* specimens were found under the bark of *Eucalyptus* sp. stone wick they clung to when it was picked up and in a reforested area in *Eucalyptus* sp., under the bark of one of these trees.

In both samples, *P. amaranthinus* were found near specimens of *Palpimanus gibbulus* Dufour, 1820 (Araneae Palpimanidae), and observing them it was noted that the way of moving of the two species was very similar. These observations linked to the similar coloring of the two species suggest some deeper co-evolutionary significance than simple causality, but this requires more careful observations and studies.

REMARKS. This record represents a new genus and a new species for Sicily and Italy.

The great adaptability of the species (Cooke, 1964) explains its wide distribution, and although it is currently present only in central southern Sicily, we are sure that future investigations will show a greater diffusion on the island.

## ACKNOWLEDGEMENTS

We want to thank our families for their constant support.

## REFERENCES

- Azevedo G.H.F, Griswold C.E. & Santos A.J., 2018. Systematics and evolution of ground spiders revisited (Araneae, Dionycha, Gnaphosidae). *Cladistics*, 34: 579–626. <https://doi.org/doi:10.1111/cla.12226>
- Bosmans R., Van Keer J., Russell-Smith A., Kronestedt T., Alderweireldt M., Bosselaers J. & De Koninck H., 2013. Spiders of Crete (Araneae). A catalogue of all currently known species from the Greek island of Crete. *Nieuwsbrief van de Belgische Arachnologische Vereniging*, 28 (Suppl. 1): 1–147
- Bosmans R., Van Keer J., Russell-Smith A., Hadjiconstantis M., Komnenov M., Bosselaers J., Huber S., McCowan D., Snazell R., Decae A., Zoumides C., Kielhorn K.-H. & Oger P., 2019. Spiders of Cyprus (Araneae). A catalogue of all currently known species from Cyprus. *Newsletter of the Belgian Arachnological Society*, 34, Suppl. 1: 1–173
- Cooke J.A.L. de, 1964. A revisionary study of some spiders of the rare family Prodidomidae. *Proceedings of the Zoological Society of London*, 142: 257–305.
- Dalmas R., 1919. Synopsis des araignées de la famille des Prodidomidae. *Annales de la Société Entomologique de France*, 87: 279–340.
- Ferrández M.A. & Carrillo J., 2018. *Prodidomus* cfr *rufus* Hentz, 1847 (Araneae: Prodidomidae), una especie “enigmática”. *Revista Ibérica de Aracnología*, 32: 123–126.
- Lucas H., 1846. Histoire naturelle des animaux articulés. In: *Exploration scientifique de l’Algérie pendant les années 1840, 1841, 1842 publiée par ordre du Gouvernement et avec le concours d’une commission académique*. Paris, Sciences physiques, Zoologie 1, 89–271, pl. 1–17.  
<https://doi.org/10.5962/bhl.title.112444>
- Nentwig W., Blick T., Bosmans R., Gloor D., Hänggi A. & Kropf C., 2021. Spiders of Europe. Version 03.2021. Online at <https://www.araneae.nmbe.ch>, accessed on 10.03.2021. <https://doi.org/10.24436/1>
- Pérez J.A. & Blasco A., 1986. Nota sobre los Prodidominae (Araneae: Gnaphosidae) de la Peninsula Iberica. *Mémoires de la Société Royal Belge d’Entomologie*, 33: 155–164
- Pickard-Cambridge O., 1872. General list of the spiders of Palestine and Syria, with descriptions of numerous new species, and characters of two new genera. *Proceedings of the Zoological Society of London*, 40: 212–354, pl. 13–16. <https://doi.org/doi:10.1111/j.1469-7998.1872.tb00489.x>
- Simon E., 1870. Sur les aranéides de la famille des Enydes qui habitent l’Espagne et le Maroc. *Revue et Magasin de Zoologie Pure et Appliquée* (2) 22: 51–54, 97–103, 142–148.
- Simon E., 1880. Description de *Diotima hirsutissima*, capture d’Araignées d’Alexandrie. *Annales de la Société Entomologique de France*, 10: 54–56.
- Simon E., 1907. Arachnides recueillis en Egypte et le long du Nil Blanc par la Mission zoologique suédoise, 1901. In: *Results of the Swedish Zoological Expedition to Egypt and the White Nile 1901 under the Direction of L. A. Jägerskiöld*. Part III. 21 (37). Uppsala, 10 pp.  
<https://doi.org/10.5962/bhl.title.21727>
- Topçu A. & Türkeş T., 2010. The first record of the family Prodidomidae (Arachnida: Araneae) in Turkey. *Turkish Journal of Zoology*, 34: 425–427
- Walckenaer C.A., 1847. Dernier Supplément. In: Walckenaer C.A. & Gervais P. (Eds.), *Histoire naturelle des Insectes. Aptères*. Tome quatrième. Roret, Paris, pp. 365–596 [Araneae: 564].  
<https://doi.org/10.5962/bhl.title.61095>
- WSC, 2021. World Spider Catalog. Version 21.5. Natural History Museum Bern, online at <http://wsc.nmbe.ch> (12.03.2021) <https://doi.org/10.24436/2>

