

Taxonomic notes on two bivalves (Mollusca Bivalvia) described by Charles-François Fontannes in 1882

Mauro M. Brunetti

Calle Navas 106, 14511 Navas del Selpillar, Spain; e-mail: mbrunetti45@gmail.com

ABSTRACT

During the study of the Pliocene malacofaunas of the Mediterranean Basin, it was ascertained that the name *Spondylus ferreolensis* Fontannes, 1882 cannot be used as a substitute for *S. concentricus* Bronn, 1831 of which the type material is here represented for the first time. The validity of *Acanthocardia perrugosa* (Fontannes, 1882) is also proposed, of which it was possible to view the type material, a species previously considered synonymous with *A. paucicostata* (GB Sowerby, 1841), or *A. bianconiana* (Cocconi, 1873) or *A. aculeata* (Linnaeus, 1758).

KEY WORDS

Pliocene; Pleistocene; Spondylidae; Cardiidae.

Received 25.10.2021; accepted 02.02.2022; published online 22.03.2022

INTRODUCTION

By re-reading the text by Fontannes (1882) we tried to clarify the taxonomy of two bivalves described by the French author. Of the first of these, *Spondylus ferreolensis* Fontannes, 1882, the comparison with the original diagnosis of *S. concentricus* Bronn, 1831 and the subsequent observations of Fontannes (1882: 210), in addition to the vision of the type material preserved at the Laboratoire de Géologie de the Université de Lyon (France), had initially suggested that *S. concentricus* could be considered *nomen dubium*. Later, however, reviewing Bronn's type material, which was thought to be lost but it is actually deposited at the Museum of Comparative Zoology at Harvard University (Cambridge), disproved this hypothesis as the two species are identical. *S. ferreolensis* must be considered as a synonym of *S. concentricus* Bronn, 1831. As for the second species, *Acanthocardia perrugosa* (Fontannes, 1882), the examination of numerous material in

addition to the syntype deposited at the Laboratoire de Géologie de l'Université de Lyon, allowed to consider the Fontannes species as a valid taxon and to speculate its paleogeographic and chronostratigraphic distribution.

MATERIAL AND METHODS

The material examined, collected during surface research, comes from various Pliocene deposits, both in the Guadalquivir basin, Spain (see Andres, 1987; Gonzales Delgado, 1985; 1988; 1989; 1993; Landau et al., 2011) and in the Zanclean/Piacenzian of Tuscany and central Lazio, Italy (Brunetti & Della Bella 2006, 2008) and southern Spain (Landau et al., 2004; 2004a; 2006; 2006a; 2007; 2011). For the generic and supra-generic determinations we followed the WoRMS (2022).

ABBREVIATIONS. La = maximum valve width; Lu = maximum length of the valve; es. =

specimen/s. LGL = Laboratoire de Géologie de Lyon, France. MNHN = Muséum National d'Histoire naturelle, Paris, France. MCZ = Museum of Comparative Zoology della Università di Harvard (Cambridge, England). CCZ = Claudio Zuccaro collection (Rome, Italy). CMB = M. Mauro Brunetti collection (Navas del Selpillar, Spain).

RESULTS

Systematics

Classis BIVALVIA Linnaeus, 1758

Superfamilia PECTINOIDEA Rafinesque, 1815

Familia SPONDYLIDAE Gray, 1826

Genus *Spondylus* Linnaeus, 1758

Type species: *Spondylus gaederopus* Linnaeus, 1758

Spondylus concentricus Bronn, 1831 (Figs. 1–6)

Spondylus concentricus Bronn, 1831: 121

Spondylus ferreolensis Fontannes, 1882: 210, table 14, figs. 3–7

Spondylus concentricus Bronn - Sacco, 1898: 6, table 3, figs. 4–8

Spondylus concentricus Bronn - Kojumdieva & Strachimirov, 1960: 75, table 26, fig. 3

Spondylus (Spondylus) concentricus (Bronn) - Malatesta, 1974: 59, table 5, figs. 1a, 1c

Spondylus concentricus Bronn - Lacour et al., 2002: 651, figs. 5F–G


Spondylus concentricus Bronn - La Croce & Repetto, 2006: p. 52, figg. 6–7.

?*Spondylus concentricus* Bronn - Chirli, 2014: 100, figs. 7–12.

MATERIAL EXAMINED. 21 es. Guidonia (Rome,



Figures 1–3. *Spondylus ferreolensis* Fontannes, 1882. Fig. 1: syntypes, Restitid (France), Lower Pliocene, with original labels, LGL. Fig. 2: syntype, inferior valve, Restitid (France), Lower Pliocene, L = 73 mm LGL. Fig. 3: syntype, inferior valve, detail of the hinge.



Figures 4–6. *Spondylus concentricus* Bronn, 1831. Fig. 4: syntypes, Bacedasco (Piacenza, Italy), Lower Pliocene, L = 70 mm, 81mm, MCZ IPBV–12458. Fig. 5: original labels. Fig. 6: Cava Formello (Guidonia, Rome, Italy) Lower/Middle Pliocene, Lu = 92 mm, CCZ.




Figure 7. *Spondylus fauroti* Jousseume, 1888, Obock, Gibuti, Upper Pliocene, syntype, Lu = 82 mm, MNHN-IM-2000-4040.

Italy), Zanclean/Piacenzian (CMB). 2 es., El Lobillo (Estepona, Malaga, Spain), Zanclean/Piacenzian (CMB).

REMARKS. Bronn (1831: 121) described *Spondylus concentricus* as a new species with this diagnosis: “*Testa ovata subobliqua; valva inferiore rugis lamelliformibus concentricis validis, in spinas validas, densas longitudinaliter seriatis productis tecta, umbone subinermi*”.

He also specifies that he only has lower valves and does not provide any type of iconography. Years later, Fontannes (1882) described *Spondylus ferreolensis* with the following original diagnosis: “*Testa ovato-oblonga, antice rotundata, postice versus cardinem paulum excavata, valde inaequalis, subtilius convexior. Valva sinistra gibbosa, irregularis, longitudinaliter costulata; costulae inaequalis, alternantes 8–9 crassioribus, primo subasperis, dein spinis subtilis*

canaliculatis munitis; umbo minimus, acuminatus, marginem cardinalem vix superans; auriculae mediocres, inaequales postica paulo majores; area cardinalis subquadrangularis, transversa; sulcus ligamenti profundus ad basin dilatatus; fossulae cardinalis subcircularis magnae, profundae; dente in extremitate rugosae vel strietae; margo palliaris valde, arcuatus, intus striatus. – Valva dextra crassior, convexior, versus area adherentum lamelli concentricis erectis munita, alibi costis longitudinalibus plus minusve obsoletis; pro partis spiniferis sculpta; area cardinalis lata, triangularis, extum paulum reflexa; margo paliaris intus denticulatus; impressio muscularis profunda, margini posteriori proximata. Dim. antero-posterior, 52; altitudo 65 mm”.

Fontannes (1882: 212) considers Bronn’s diagnosis, lacking a figures, nomen dubium, since, in his opinion, there is no particular character of a distinct species. Bronn’s type material (Figs. 4, 5) is identi-

cal to the Fontannes species and shows this hypothesis was wrong. *S. ferreolensis*, the illustrated type series (Figs. 1–3), is therefore to be considered synonymous with *S. concentricus* (Strachimirov, 1960; Lacour et al., 2002). This taxon, which is poorly reported in literature, is known with certainty only from the upper Miocene onwards. Markedly thermophilic species, it becomes extinct during the Piacenzian period. According to Malatesta (1974), it is found only in clayey sediments, as other more recent findings confirm (Fig. 6). The same author considered as a synonymous, in my opinion erroneously, *S. fauroti* Jousseume, 1888 (Fig. 7), from the Upper Pleistocene of Djibouti, with *S. concentricus*, from which it differs at first glance, for its significantly more regular sculpture of the valve inferior. The specimen figured by Chirli (2014) does not seem to correspond to *S. concentricus* as it lacks its characteristic deep ligamentary sulcus (Fig. 3).

Superfamilia CARDIOIDEA Lamarck, 1809

Familia CARDIIDAE Lamarck, 1809

Genus *Acanthocardia* Gray, 1851

Type species: *Acanthocardia aculeata* (Linnaeus, 1758)

Acanthocardia perrugosa (Fontannes, 1882) (Fig. 8–11)


Acanthocardia aculeata var. *perrugosa* Fontannes, 1882: 81, table 5, figs. 2–3

Cardium paucicostatum Sowerby - Dollfus et al., 1904: 43, table 15, figs. 6–7

Cardium ciliare bianconianum (Cocconi) - Glibert & Van de Poel, 1970: 55

Acanthocardia perrugosa Fontannes - Domench, 1984: 10

Acanthocardia (*A.*) *perrugosa* (Fontannes) - Martinell & Domench, 1984: 12, table 5, figs. 3–4



Figures 8–10. *Acanthocardia perrugosa* (Fontannes, 1882). Figs. 8, 9: Syntypes, Lower Pliocene, La = 42 mm, LGL–EM353380. Fig. 10: juvenile specimen, Lucena del Puerto (Huelva, Spagna), Lower Pliocene, La = 27 mm CMB.



Figures 11–13. *Acanthocardia* spp. Fig. 11: *Acanthocardia perrugosa* (Fontannes, 1882), Lucena del Puerto (Huelva, Spain) Lower Pliocene, La = 51 mm, CMB. Fig. 12: *Acanthocardia paucicostata* (G.B. SowerbyII; 1834), Villalvernia (Alessandria), Lower Pliocene, La = 50.8 mm CMB. Fig. 13. *Acanthocardia bianconiana* (Cocconi, 1873), Torrente Arda (Piacenza), Calabrian, La = 51.5 mm CMB.



Figures 14–16. *Acanthocardia* spp. Fig. 14: *Acanthocardia echinata* (Linnaeus, 1758), Cutrofiano (Lecce), Calabrian, La = 45 mm CMB. Figs. 15, 16: *Acanthocardia paucicostata* (G.B. Sowerby II, 1834). Fig. 15: Torrente Enza (Reggio Emilia), Gelasian, La = 24 mm CMB. Fig. 16: Mazagon (Huelva, Spain), Recent, La = 30 mm.

Acanthocardia (*Acanthocardia*) *paucicostata* “morfofoto tipo *bianconianum* Cocconi” - Andrés, 1987: 115, table 3, figs. 5–7

Acanthocardia (*Acanthocardia*) *paucicostata* Sowerby - Lozano Francisco, 1997, table 44, figs. 1, 2.

Acanthocardia paucicostata Sowerby - Cárdenas et al., 2017: 376, fig. 7J

Acanthocardia aculeata (Linnaeus, 1758) - Pimental, 2018, table 8, figs. 5–6

MATERIAL EXAMINED. 22 es. Lucena del Puerto (Huelva, Spain), Zanclean (CMB). 25 es. Santa Catalina (Huelva, Spain), Zanclean (CMB). 10 es. El Lobillo (Estepona, Malaga, Spain), Zanclean/Piacenzian (CMB). 4 es. Villarsa (Huelva, Spagna), Zanclean (CMB).

ORIGINAL DESCRIPTION (Fontannes, 1882): “*Testa minor, minus obliqua postice minus truncata, umbones plerumque tumidiores; costae 17–18 subtriangulares in medio carinatae; rugae interstitiorum crassiores, irregulares, super costae sine alteratione transeuntes. Diam. antero-post., 42; altitudo, 40 millim*”.

REMARKS. Morphological characteristics of this species are the relatively low number of ribs (17–18), the slightly oblique shape, the swollen umbo, the triangular profile ribs crossed by the growth lines and equipped with sturdy spines, short, in the juvenile specimens long and thin. *Acanthocardia perrugosa*, often misinterpreted, has been considered time and again as a variety of *A. paucicostata* (Sacco, 1898; Dollfus et al., 1904) of *A. bianconiana* (Glibert & Van de Poel, 1970; Andrés, 1987)

or *A. aculeata* (Pimental, 2018). To understand how this could have happened, it is necessary to consider two facts. First of all, the Fontannes species, as here discussed, appears to be present only in the Tortonian of Portugal (Dollfus et al., 1904) and Spain (Cárdenas et al., 2017), in the Lower Pliocene in the westernmost part of the Mediterranean basin, from southern France to Portugal and in the Guadalquivir basin, where it is found very common, reaching considerable dimensions (>70 mm, pers. obs.), while it is absent in the Italian Pliocene and the eastern Mediterranean. Thus, for example, the exemplars figured by Sacco (1898), of the Italian Pliocene, which he interpreted as *A. perrugosa*, all belong to either *A. paucicostata* or *A. bianconiana*. The second observation is that the populations of *A. paucicostata* of the Lower/middle Pliocene (Fig. 12) are often larger than the current populations, including Atlantic ones, with fewer ribs and with other ribs that are slightly more prominent than spines. This may have induced some authors to confusion. The forms of *A. paucicostata* close to the current ones (Fig. 15) begin to appear only starting from the Gelasian (pers. obs.). The juvenile specimens of *A. perrugosa* with thorniness (Fig. 13) may have misled Fontannes who described *A. perrugosa* as a variety of *A. aculeata* species with a different, sub-quadrangular shell shape and with a greater number of ribs and a consequent more “thorny” aspect. La Perna & D’Abramo (2013) clarified the differences between *A. paucicostata* and *A. bianconiana* very well, while they are doubtful about the validity or otherwise of *A. perrugosa*. I do not agree with some statements of these two authors, first of all in considering fig. 2 and fig. 3 of Fontannes as two different species, presenting the same number of ribs and similar sculpture. Furthermore, the fact that he considered Coppi and not Cocconi the author of *A. bianconiana*, a species that he knew well, could simply be a *lapsus calami*. The specimen illustrated by Martinell & Domenich (1984, plate 5, figs. 3, 4) is confirmed to belong to *A. perrugosa*, contrary to what is stated by La Perna & D’Abramo (2013).

ACKNOWLEDGEMENTS

Thanks are due to Emmanuel Robert, CNRS, Research Engineer, PhD, Geosciences Collections

Manager Laboratoire de Géologie de Lyon (France) for the photos of the Fontannes types; to Jessica D. Cundiff, Curatorial Associate, Department of Invertebrate Paleontology, Museum of Comparative Zoology Harvard University (Cambridge, England) for the photos of Bronn’s types; to my friends: Ermanno Quaggiotto (Longare, Vicenza) for bibliographic help, Claudio Zuccaro (Guidonia, Rome, Italy) for the numerous material he made available, Massimo Cresti (San Casciano VP, Florence, Italy) for proofreading the text and giving some valuable suggestions.

REFERENCES

- Andrés I., 1987. Los Heterodonta (Bivalvia) en el Plioceno marino de Bonares (Huelva). *Studia Geologica Salmanticensia*, 24: 83–149.
- Bronn H.G., 1831. Italiens Tertiär-Gebilde und deren organische Einschlüsse. K. Groos, Heidelberg.
- Brunetti M.M. & Della Bella G., 2006. *Leufroyia ferrierii*: una nuova specie per il Pliocene toscano (Gastropoda: Conidae). *Bollettino Malacologico*, 42: 118–120.
- Brunetti M.M. & Della Bella G., 2008. *Macalia* (?) *kengii* n. sp. un nuovo bivalve per il Pliocene toscano (Tellinidae, Macominae). *Bollettino Malacologico*, 44: 115–118.
- Cárdenas J., Bajo I. & Maestre M.V., 2017. Estudio paleontológico de los bivalvos (Mollusca) del Tortonense superior de Arroyo Trujillo, Cantillana (Sevilla). *Spanish Journal of Palaeontology*, 32: 367–386.
- Chirli C., 2014. Malacofauna pliocenica toscana vol. 10 Bivalvia Pteriomorpha Beurlen, 1884. Tipografia Vanzì S.r.l., Colle Val d’Elsa, Siena, 220 pp.
- Dollfus G.F., Berkeley Cotter J.C. & Gomes J.P., 1903–1904. Planches de Céphalopodes, Gastéropodes et Pélécypodes laissée par F.A. Pereira da Costa accompagnée d’une explication sommaire et d’une esquisse géologique. Imprimerie de l’Académie Royale des Sciences, Lisbonne, 206 pp.
- Glibert M. & van de Poel L., 1970. Les Bivalvia fossiles du Cénozoïque étranger. *Memoires de l’Institut royal des Sciences naturelles de Belgique*, 2° serie, 84: 1–185.
- Gonzales Delgado J.A., 1985. Estudio sistemático de los Gastéropodos del Plioceno de Huelva (SW de España). 1. Archeogastropoda. *Studia Geologica Salmanticensia*, 20: 45–77.
- Gonzales Delgado J.A., 1988. Estudio sistemático de los Gastéropodos del Plioceno de Huelva (SW de España). 3. Mesogastropoda (Scalacea-Tonnacea). *Studia Geologica Salmanticensia*, 25: 109–160.

- Gonzales Delgado J.A., 1989. Estudio sistématico de los Gastéropodos del Plioceno de Huelva (SW de España). 3. Neogastropoda (Muricea-Buccinacea). *Studia Geologica Salmanticensia*, 26: 269–315.
- Gonzales Delgado J.A., 1993. Estudio sistématico de los Gastéropodos del Plioceno de Huelva (SW España). 5. Neogastropoda (Volutacea - Conacea). *Studia Geologica Salmanticensia*, 28: 7–69.
- Kojumdgieva Em. & Strachimirov B., 1960. Les Fossiles de Bulgarie, VI Tortonien. Sofia, Academie des Sciences de Bulgarie, 317 pp.
- Lacour D., Lauriat-Rage A., Saint Martin J.-P., Videt B., Néraudeau D., Goubert E. & Bongrain M., 2002. Les associations de bivalves (*Mollusca, Bivalvia*) du Messinien du bassin de Sorbas (SE Espagne). In: Néraudeau D. & Goubert E. (Eds.), *l'Événement messinien: approches paléobiologiques et paléocéologiques*. *Geodiversitas*, 24: 641–657.
- Lacroce L. & Repetto C., 2003. La famiglia Spondylidae Gray, 1826 nel Pliocene italiano. *La Conchiglia*, 306: 48–54.
- Landau B., Da Silva C.M. & Mayoral E., 2011. The Lower Pliocene gastropods of the Huelva Sands Formation, Guadalquivir Basin, Southwestern Spain. *Palaeofocus*, 4: 1–90.
- Landau B., Marquet R. & Grigis M., 2003. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 1. Vetigastropoda. *Palaeontos*, 3: 1–87.
- Landau B., Marquet R. & Grigis M., 2004. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 2. Orthogastropoda, Neotaenioglossa. *Palaeontos*, 4: 1–108.
- Landau B. & Feshe D., 2004a. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 3. Trivioidea, Cypraeoidea. *Palaeontos*, 5: 1–34.
- Landau B., Beu A. & Marquet R., 2004a. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 5. *Palaeontos*, 5: 35–102.
- Landau B. & Da Silva C.M., 2006. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 9: Olividae. *Palaeontos*, 9: 1–21.
- Landau B., La Perna R. & Marquet R., 2006a. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 10: Marginellidae, Cystiscidae. *Palaeontos*, 9: 22–60.
- Landau B., Petit R. & Marquet R., 2006. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 12: Cancellarioidea. *Palaeontos*, 9: 61–101.
- Landau B., La Perna R. & Marquet R., 2006a. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 6: Triphoroidea, Epitonioidea, Eulimoidea. *Palaeontos*, 10: 1–96.
- Landau B., Houart R. & Marquet R., 2007. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 7: Muricidae. *Palaeontos*, 11: 1–87.
- Landau B., Da Silva C.M. & Gili C., 2009. The Early Pliocene Gastropoda (*Mollusca*) of Estepona, southern Spain. Part 8: Nassariidae. *Palaeontos*, 17: 1–101.
- Landau B., Da Silva C.M. & Mayoral E., 2011. The Lower Pliocene gastropods of the Huelva Sands Formation, Guadalquivir Basin, Southwestern Spain. *Palaeofocus*, 4: 1–90.
- Lozano-Francisco M.C., 1997. Los bivalvos del Plioceno de la provincia de Málaga. Tesis Doctoral Depto. Ecología y Geología. Facultad de Ciencias. Universidad de Málaga, 1016 pp.
- Malatesta A., 1974. *Malacofauna pliocenica umbra. Memorie per servire alla descrizione della carta Geologica Italiana Servizio Geologico d'Italia*. Roma, Tipografia Pinto, 498 pp.
- Martinell J. & Domènech R., 1984. *Malacofauna del Plioceno de Sant Onofre (Baix Ebre; Tarragona)*. *Iberus*, 4: 1–27.
- Sacco F., 1898. *I Molluschi dei terreni terziari del Piemonte e della Liguria*. Torino, Ed. Clausen, vol. 26, 96 pp.
- WoRMS, 2022. <http://www.marinespecies.org/index.php> del 1/1/2021