

# The unusual occurrence of *Chaetomorpha antennina* (Bory) Kützinger, 1847 (Chlorophyta Cladophoraceae) in a sublittoral rocky reef of the Mexican tropical Pacific marine ecoregion: significant expansion of their ecogeographic range

Norma López\* & Carlos Candelaria-Silva

Escuela Nacional de Estudios Superiores Morelia. Universidad Nacional Autónoma de México. Antigua Carretera a Pátzcuaro No. 8701, Col. Ex Hacienda de San José de la Huerta, C.P. 58190 Morelia, Michoacán, México

\*Corresponding author, e-mail: [nlopez@enesmorelia.unam.mx](mailto:nlopez@enesmorelia.unam.mx)

## ABSTRACT

This is the first record of *Chaetomorpha antennina* (Bory) Kützinger, 1847 (Chlorophyta Cladophoraceae) from a deep rocky reef of the Mexican tropical Pacific marine ecoregion, a chlorophyte which had been reported only from intertidal and shallow subtidal zones. *C. antennina* of Sacramento reef in Zihuatanejo, Mexico, is smaller than had been previously reported. Extension of variation intervals of this species collected 18–20 m depth suggests that other factors such as hydrodynamic forces and light reduction in deep environments may have a relevant role in its morphological plasticity.

## KEY WORDS

*Chaetomorpha antennina*; subtidal algae; deep rocky reef; Eastern tropical Pacific; Zihuatanejo.

Received 11.05.2022; accepted 22.09.2022; published online 24.10.2022

## INTRODUCTION

The phycofloristic knowledge of the Mexican tropical Pacific marine ecoregion (Spalding et al., 2007) has increased notably since the 1980s and currently there is a clear picture of the presence and distribution of the majority of the reported species (González-González, 1992; Pedroche & Senties, 2003). Such is the case for the chlorophyte *Chaetomorpha antennina* (Bory) Kützinger 1847 (Phylum Chlorophyta, Order Cladophorales), which has a thallus with a simple uniseriate level of filamentous organization, forming upright bushes constituted by aggregations of numerous filaments [see photographs in Guiry (2022)]. It is an epilithic species with a pantropical distribution; it is one of the most prominent species in the Mexican tropical Pacific

and it is well known ecologically. It has been reported in numerous locations at all points along the 1,200 kilometers of the coast (Pedroche et al., 2005; León-Álvarez, 2016) in different rocky habitats and in a wide range of microenvironmental conditions (González-González, 1993; Candelaria et al., 2006; Norris, 2010). This species is commonly found from the high intertidal zone to the low intertidal zone (Norris 2010; Leliaert et al., 2011a) and very rarely in the subtidal shallows to a depth of 2 meters (Mendoza-González et al., 2018). It can be found on more or less vertical cliff walls, rocky points, rocky blocks or breakwaters. It is generally directly exposed to the swell, lateral dragging, splash or intense spray. It is exposed to conditions of direct illumination, with more or less prolonged periods of sunshine.

Although it is true that there is already an extensive information base regarding the phycoflora of the Mexican tropical Pacific, there is a notable bias toward intertidal algae and comparatively little is known regarding sublittoral algal biodiversity. As part of a project focused on the algae of the rocky reefs in the Zihuatanejo area, in the state of Guerrero, we collected *C. antennina* at a depth of 18–20 meters. Given that this finding is very significant in terms of the ecogeographical distribution of this species, the objective of this study is to provide the first report of *C. antennina* in an unusual habitat in the marine ecoregion of the Mexican tropical Pacific.




Figure 1. Location of the state of Guerrero, southern Mexico (upper) and Sacramento rocky reef in the Ixtapa-Zihuatanejo region (lower).

## MATERIAL AND METHODS

We carried out a survey sampling in the Sacramento rocky reef (17°38'02"N, 101°36'39"W), at southern Ixtapa-Zihuatanejo (Fig. 1). This site descends to a depth of 30 m and is composed of overlapping, irregularly shaped rocky blocks 1–5 m long, among which there are deposits of sand, gravel and boulders, which are exposed to strong internal currents that produce laminar or turbulent flows of water. *Chaetomorpha antennina* is attached to rocks and balanus shells.

The samples were collected randomly by SCUBA diving on 23 October 2012 between 18 and 20 m depth. All the algal specimens were collected by hand using hammer and chisel and preserved in 4% formalin in seawater for morphological analysis. The specimens were deposited in Sección de Algas, Herbario de la Facultad de Ciencias (FCME), Universidad Nacional Autónoma de México.

The morphological characteristics were examined following taxonomic descriptions of *C. antennina* from the Mexican tropical Pacific (Setchell & Gardner, 1924; Taylor, 1945; Abbott & Hollenberg, 1979; Candelaria, 1985; López, 1993; Norris, 2010; Leliaert et al., 2011a). Diagnostic characters used were: habit, thallus length, basal cell features (shape, length, upper diameter, presence of annular constrictions and type of holdfast), suprabasal cell length, and intermediate cells features (shape, length and diameter). Morphometric analyses were carried out with an Olympus SZ2ILST stereo microscope and an Olympus BX51TF microscope. Photographs were taken with an Olympus SC100 digital camera mounted on an Olympus SZX10 stereomicroscope and Olympus cellSens Standard software.

## RESULTS

The recognition of *Chaetomorpha* species is currently based on a few morphological features such as growth form, cell shape and dimensions, and shape of basal attachment cell (Leliaert et al. 2011a, 2011b). The collected specimens form tufts of scarce fine green filaments, 0.6–1.2 cm high (Fig. 2). The elongated, thick-walled basal cells are club-shaped, 600–3600 µm long and 180–570 µm in diameter at


the distal end; with tenuous annular constrictions near the base; proximal pole bearing branched rhizoids (Fig. 3). The suprabasal cells are 185–720  $\mu\text{m}$  long (Fig. 4). Other cells are sub cylindrical (Fig. 5) towards the basal area of the filaments, 510–900  $\mu\text{m}$  long and 360–840  $\mu\text{m}$  in diameter, gradually becoming broader and barrel-shaped upward (Fig. 6). Reproductive cells were not observed.

Sacramento specimens are in accordance with *C. antennina* descriptions from the Mexican tropical Pacific in habit and length of thallus; form, length and diameter of basal cell; presence of annular constrictions; presence of branched rhizoids; suprabasal cell length; and shape, length and diameter of intermediate cells. However, other features, specifically measurements, extend down the recorded variation intervals, e.g. thallus length (our specimens 0.6–1.2 cm vs. 1–16 cm from previous reports); basal cell length (600–3600  $\mu\text{m}$  vs. 1800–15000  $\mu\text{m}$ ); suprabasal cell length (185–720  $\mu\text{m}$  vs. 360–3700  $\mu\text{m}$ ) (Table 1). Also, the scarcity of filaments in tufts in our material was notable, in contrast to the dense bush described by numerous

authors from many Mexican tropical Pacific localities. Furthermore, the presence of annular constrictions is barely distinguishable.

## DISCUSSION

*Chaetomorpha antennina* has been recognized as a species that is “clearly circumscribed and shows a fairly constant morphology throughout its ecological and geographical range” (Leliart et al., 2011a) and it is easily identifiable from other species in the genus essentially from its habit, a long claviform basal cell, branched rhizoids that develop on the proximal extreme of the basal cell, and the presence of annular constrictions. In turn, Abbott and Huisman (2004) mention that “the identification of *C. antennina* is confirmed by the presence of a conspicuously long basal cell”, just as occurs in our specimens. Considering the morphologic definition of this species, there is no doubt about the identity of *C. antennina* from Sacramento reef, however the habit formed by sparse filaments



Figures 2-6. *Chaetomorpha antennina* from Sacramento Reef. Fig. 2: a fine filament. Fig. 3: branched rhizoids. Fig. 4: suprabasal cell. Fig. 5: intermediate cells. Fig. 6: apical cell showing upper end. Scale bars: Figs. 2, 5 = 500  $\mu\text{m}$ ; Fig. 3, 6 = 200  $\mu\text{m}$ ; Fig. 4 = 450  $\mu\text{m}$ .

and the smaller thallus length were noteworthy. This aspect represents a pattern of size reduction similar to that observed in many subtidal turf-forming species from Zihuatanejo (López et al., 2004). Also noteworthy, is the barely distinguishable presence of annular constrictions on the basal cell. Several authors indicate that this characteristic is commonly (Taylor, 1945), frequently (Abbott & Hollenberg, 1979), or usually (Norris, 2010) present; however, Leliart et al. (2011a) mention that in culture conditions, the annular constrictions do not appear, which could be associated with a morphogenetic development in conditions of lower hydrodynamics in the sublittoral or the non-existent movement in cultures, when compared with the prevailing conditions in the intertidal zone.

The impact that abiotic factors have on deep habitat organisms is considerably different from that which occurs on organisms of intertidal habitats

(Coleman, 2002; Araújo et al., 2005; Benedetti-Cecchi & Trussell, 2014) or subtidal shallow rocky habitats (Garrahou et al., 2002; Balata & Piazzini, 2008). The tides, swell, desiccation, light, temperature, and nutrient availability, are among the most extreme and dynamic factors that *C. antennina* is exposed to in the intertidal zone. The differences found in *C. antennina* from the Sacramento reef could be interpreted as a plastic response to the environmental gradients that are relatively more constant in deep habitats, such as the low incidence of light, lower temperatures, and lower intensity turbulence from water currents, similar to the conditions observed in deep rocky reefs in the Mediterranean (Garrahou et al., 2002; Balata & Piazzini, 2008), but it is necessary to evaluate this correlation at Sacramento and at other rocky reefs in the Mexican tropical Pacific. The ecological conditions in the rocky habitats of the intertidal and sub-

Features	Previously reported features	Our description
Habit	Erect brushlike tufts	Tufts of very fine green scarcity filaments
Length of thallus	1-16 cm	0.6-1.2 cm
Basal cell form	Club-shaped	Club-shaped
Basal cell length	1800-15000 µm	600-3600 µm
Basal cell diameter	180-900 µm	180-570 µm
Annular constrictions	Conspicuous annular constrictions generally	Tenuous annular constrictions
Presence of branched rhizoids	Proximal end bearing branched rhizoids	Proximal end bearing branched rhizoids
Suprabasal cell length	360-3700 µm	185-720 µm
Intermediate cell form	Subcylindrical towards the basal area of the filaments, gradually becoming broader and barrel-shaped upward	Subcylindrical towards the basal area of the filaments, gradually becoming broader and barrel-shaped upward
Intermediate cell length	330-2000 µm	510-900 µm
Intermediate cell diameter	225-1000 µm	360-840 µm

Table 1. Features of *Chaetomorpha antennina* from the Sacramento rocky reef and those previously reported.

tidal shallows where *C. antennina* has been previously reported and those which we are reporting at the deep rocky reef of Sacramento, suggest that this species has an adaptive capacity greater than was previously known in this ecoregion and expands its condition as an eurytopic species (Lincoln et al., 2001), exhibiting a tolerance for an ample variety of habitats, from the upper intertidal zone to the subtidal depths (18–20 m), and therefore its ecological range is extended as has been shown in this work.

This report is a contribution to morphological and ecogeographical knowledge of *Chaetomorpha antennina* in the Mexican tropical Pacific, and highlights the need to carry out more detailed studies of distribution patterns, habitat characterization, and morphological descriptions of subtidal flora, which constitute the baseline of rocky reefs biodiversity of this ecoregion, as was pointed out by Rincón-Díaz et al. (2020) for the Panama Bight ecoregion at the Eastern Tropical Pacific.

## ACKNOWLEDGEMENTS

We thank M. Sc. Pedro Ramírez-García for his assistance in the field, Dr. César Andrés Torres-Miranda and M. Sc. Douglas Castillejos-Lemus for their advice during photographic sessions, Alejandra Cruz-Rodríguez for image editing, David Silva for the map and Josué Lazcano for his support in consulting the Herbarium's information system. The authors acknowledge the infrastructure provided by Escuela Nacional de Estudios Superiores, Morelia, Universidad Nacional Autónoma de México. We also thank M. Sc. Michele Gold-Morgan and Nick Wolf for their valuable assistance with the English version.

## REFERENCES

- Abbott I.A. & Hollenberg G.J., 1974. Marine Algae of California. Stanford University Press, Stanford, 827 pp.
- Abbott I.A. & Huisman, J.M., 2004. Marine green and brown algae of the Hawaiian Islands. Bishop Museum Press, Honolulu, 259 pp.
- Araújo R., Bárbara I., Sousa-Pinto I. & Quintino V., 2005. Spatial variability of intertidal rocky shore assemblages in the northwest coast of Portugal. Estuarine, Coastal and Shelf Science, 64: 658–670. <https://doi.org/10.1016/j.ecss.2005.03.020>
- Balata D. & Piazzì L., 2008. Patterns of diversity in rocky subtidal macroalgal assemblages in relation to depth. Botanica Marina, 51: 454–471.
- Benedetti-Cecchi L. & Trussell G.C., 2014. Intertidal Rocky Shores. In: Bertness M.D., Bruno J.F., Silliman B.R. & Stachowicz J.J., Marine Community Ecology and Conservation. Sinauer Associates, Inc., Oxford, pp. 203–225.
- Candelaria C., 1985. Caracterización de la ficoflora de la localidad de Puerto Escondido, Guerrero. Bachelor dissertation, Facultad de Ciencias, Universidad Nacional Autónoma de México, 77 pp.
- Candelaria C., Rodríguez D., López N. & González-González J., 2006. Patrón de distribución de macroalgas en un canal de corrientes. TIP Revista Especializada en Ciencias Químico-Biológicas (México), 9: 65–72.
- Coleman M.A., 2002. Small scale spatial variability in intertidal and subtidal turfing algal assemblages and the temporal generality of these patterns. Journal of Experimental Marine Biology and Ecology, 267: 53–74. [https://doi.org/10.1016/S0022-0981\(01\)00358-6](https://doi.org/10.1016/S0022-0981(01)00358-6)
- Garrabou J., Ballesteros E. & Zabala M., 2002. Structure and dynamics of North-Western Mediterranean Rocky Benthic Communities along a Depth Gradient. Estuarine, Coastal and Shelf Science, 55: 493–508.
- González-González J., 1992. Estudio florístico ecológico de ambientes y comunidades algales del litoral rocoso del Pacífico Tropical Mexicano. PhD dissertation, Facultad de Ciencias, Universidad Nacional Autónoma de México, 167 pp.
- González-González J., 1993. Comunidades algales del Pacífico Tropical. In: Salazar-Vallejo S.I. & González N.E. 1993. Biodiversidad Marina y Costera de México. CONABIO & CIQRO, México, 420–443.
- Guiry M.D., in Guiry M.D. & Guiry G.M. 2 de marzo de 2021. AlgaeBase. World-wide electronic publication, National University of Ireland, Galway. <https://www.algaebase.org>; searched on April 29th, 2022.
- Leliaert F., D'hondt S., Tyberghein L., Verbruggen H. & De Clerck O., 2011a. Atypical development of *Chaetomorpha antennina* in culture (Cladophorales, Chlorophyta). Phycological Research, 59: 91–97. <https://doi.org/10.1111/j.1440-1835.2010.00604.x>
- Leliaert F., Payo D.A., Calumpong H.P. & De Clerck O., 2011b. *Chaetomorpha philippinensis* (Cladophorales, Chlorophyta), a new marine microfilamentous green alga from tropical waters. Phycologia, 50: 384–391. <https://doi.org/10.2216/10-76.1>
- Lincoln R.J., Boxshall G.A. & Clark P.F., 2001. A Dictionary of Ecology, Evolution and Systematics. Cambridge University Press, Cambridge, 360 pp.



- López N., 1993. Caracterización de la ficoflora sublitoral de Acapulco y Zihuatanejo, Gro. Bachelor dissertation, Facultad de Ciencias, Universidad Nacional Autónoma de México, 89 pp.
- López N., Rodríguez D. & Candelaria C., 2004. Intraspecific morphological variation in turf-forming algal species. *Universidad y Ciencia, Special Num.*, 1: 7–15.
- Mendoza-González A.C., Mateo-Cid L.E., Alvarado-Villanueva R., Sotelo-Cuevas F., Ceballos-Corona J.G.A. & Garduño-Acosta A.G.A., 2018. Nuevos registros y lista actualizada de las algas verdes (Chlorophyta) del litoral de Michoacán, México. *Revista Mexicana de Biodiversidad*, 89: 971–985.  
<https://doi.org/10.22201/ib.20078706e.2018.4.2604>
- Norris J.N., 2010. Marine Algae of the Northern Gulf of California: Chlorophyta and Phaeophyceae. *Smithsonian Contributions to Botany*, 94: 1–276.
- Pedroche F.F. & Senties A., 2003. Ficología marina mexicana. *Diversidad y Problemática actual. Hidrobiológica*, 13: 23–32.  
<https://hidrobiologica.izt.uam.mx/index.php/revHidro/article/view/1093>
- Pedroche F.F., Silva P.C., Aguilar-Rosas L.E., Dreckmann K.M. & Aguilar-Rosas E., 2005. Catálogo de las algas marinas bentónicas del Pacífico de México. I. Chlorophycota. Universidad Autónoma Metropolitana-Iztapalapa/University of California/Universidad Autónoma de Baja California, México, 136 pp.
- Rincón-Díaz N., Gavio B., Sánchez Muñoz J.V. & Chasqui L., 2020. *Crouania mageshimensis* Itono, 1977 (Ceramiales, Rhodophyta) and new records for three other species of macroalgae from the Eastern Tropical Pacific. *Check List*, 16: 1171–1180.  
<https://doi.org/10.15560/16.5.1171>
- Setchell W. & Gardner N.L., 1920. The marine algae of the Pacific coast of North America. Part II. Chlorophyceae. *University of California Publications in Botany*, 8: 139–374.
- Spalding M.D., Fox H.E., Allen G.R., Davidson N., Ferdaña Z.A., Finlayson M., Halpern B.S., Jorge M.A., Lombana A., Lourie S.A., Martin K.D., McManus E., Molnar J., Recchia C.A. & Robertson J., 2007. Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *BioScience*, 57: 573–583.  
<https://doi.org/10.1641/B570707>
- Taylor W.R., 1945. Pacific marine algae of the Allan Hancock Expeditions to the Galapagos Islands. *Allan Hancock Pacific Expeditions*, 12: 1–528.