

Eco-biological study of the Largemouth bass, *Micropterus salmoides* Lacépède, 1802 (Perciformes Centrarchidae), from Keddara Dam (Boumerdes, Algeria)

Chabet dis Chalabia^{1,2*}, Ferhani Khadra¹, Laababsa Leila¹, El Houatti Habiba¹, Zouaoui Karima³, Boucena Mohamed Amine¹, Itchir Rachida¹, Chelif Halim¹, Belhouchet Ismail², Redjemi Momahed Aimen² & Didani Amira¹

¹Centre National de Recherche et de Développement de la Pêche et de l'Aquaculture (CNRDPA), Boulevard front de mer, Bou Ismaïl 42415 Tipaza, Algeria

²École Nationale Supérieure des Sciences de la Mer et de l'Aménagement du Littoral (ENSSMAL), Campus Universitaire de Dely Ibrahim Bois des Cars 'Dély Ibrahim 16320, Algiers, Algeria

³Université Saad Dahleb, Blida 1, Faculté des Sciences de la Nature et de Vie, Département des Biotechnologies, Blida, Algeria

*Corresponding author: ch.d.chalabia@gmail.com

ABSTRACT

The aim of this work is to provide necessary information on the eco-biology of Largemouth bass *Micropterus salmoides* Lacépède, 1802 (Perciformes Centrarchidae) in Keddara Dam Lake, located in northern part of Algeria. A total of 67 specimens (22 females and 45 males) were collected during December 2019 and September 2020. The sex ratio was M:F = 2.33:1 and M:F = 2:1 for December 2019 and September 2020, respectively. The length-weight relationship was estimated for each period, with $W = 0.1672 \times L^{2.291}$ for December 2019 and $W = 0.0039 \times L^{3.405}$ for September 2020. The condition factor K estimated for December 2019 was $K = 15.51 \pm 0.49$, for September 2020 was $K = 13.92 \pm 2.22$. Macroscopic observation of the gonads showed that the spawning period begins in December. *Micropterus salmoides* feeding activity was high in December 2019, mostly eating fish (60% and 24.56% for fish part for December 2019 and September 2020, respectively, and 10% and 26.31% for full fish for December 2019 and September 2020, respectively). Largemouth bass display biological adaptation and trophic plasticity possibly facilitating its success in Keddara Dam as invasive species.

KEY WORDS

Feeding; gonads; invasive; length-weight; sex ratio.

Received 27.11.2022; accepted 24.04.2023; published online 10.08.2023

INTRODUCTION

Nine perch-like freshwater fish were farmed in 30 countries worldwide, with a total production of 0.8 million tonnes, with worth of around US\$ 7 billion (Cai et al., 2019), accounting for approximately 0.7% of total tonnage (McLean et al., 2022). Largemouth bass, *Micropterus salmoides* Lacépède, 1802

(Perciformes Centrarchidae, is in the first place in this group, representing for 56 percent of the total production (Cai et al., 2019). China accounts the vast majority of Largemouth bass production (Hussein et al., 2020), which stood at 432.000 tonnes in 2018 (CFS, 2019).

Largemouth bass *Micropterus salmoides*, is an important species in recreational fisheries in North

America since the 1880s and later became the most widely popular and distributed gamefish in the United States since 1975 (Cooke & Philipp, 2009). This species have been introduced in Algeria for the first time in 1956 by the French for the recreational fisheries (Kara, 2012). Beyond some information on spawning and reproduction under controlled conditions (CNRDPA, unpublished data; Zouakh & Meddour, 2017), little is known of the ecology and biology of Largemouth bass in its natural habitat. Lakes and dams may furthermore represent nursery areas for Largemouth bass (CNRDPA, unpublished data), which could yield insights into conditions that affect the recruitment of this species.

Micropterus salmoides has become an economically important freshwater aquaculture species in Algeria due to several advantages including good flesh quality, short culture cycle, rapid growth performance, tolerance to handling, and strong adaptability (Bai & Li, 2019).

Determinations of ecological and biological characteristics are fundamental in understanding the adaptation success of Largemouth bass in Algeria biotope, as they allow for assessment of the establishment success of this species, reporting paucity of information on biology and ecology of this species introduced range in Africa (Azuma & Motomura, 1998; Britton et al., 2010; Taylor & Weyl, 2017) and the scarcity of data on its biology and ecology in Algeria.

The present study aimed to provide information and improve the understanding of the adaptation success of Largemouth bass by assessing reproduction, growth, sex ratio, and diet in *Micropterus salmoides* populations from the Keddara Dam in Northeast Algeria. We report for the first time this species in this lake and, using the results from capture in two different periods (beginning and end of the autumn season), we studied this species in relation to growth, reproduction, and diet and its environment.

MATERIAL AND METHODS

Study area

The study was carried out in Keddara Dam (36°65'N; 3°43'E), one of the most important dams in Algeria's central region. The reservoir has a total

surface area of 5.2 km² and a maximum depth of 150 m.

Conductivity (µs/cm), turbidity (NTU), pH, and salinity (PSU) of Keddara Dam were measured in situ using a multi-parameter analyzer (Model WTW multi 340i) and transparency using a Secchi disk.

Material

Micropterus salmoides is caught with a trammel net of 50 mm mesh size, 200 m in length, and 3.5 m in width, during December 2019 and September 2020. Immediately after landing, we recorded for each fish total length (cm), standard length (cm), fork length (cm), total weight (g), gutted weight (g), and weights of the liver (g), gonad (g), and stomach (g). Sex was determined by macroscopic examination of the gonads according to stages established by Holden & Raitt (1974) of which we find five stages of sexual maturity for both males and females (stage I: immature; stage II: maturing virgin and recovering spent; stage III: pre-spawning; stage IV: spawning; stage V: spent/post-spawning phase). The Fulton's condition factor, $K = (\text{body weight (g)} / \text{total length (cm)}^3) * 100$, were calculated for both sexes. The length-weight relationships of Largemouth bass were also estimated separately for each autumn (2019–2020).

The stomach contents analysis was made after storage at 5% formalin. The identification of the prey was conducted at a higher level of taxonomic rank (subphylum). For the diet analysis, vacuity index (VI), frequency of occurrence (O), numerical abundance (N), and predation intensity (PI), were calculated as (Hynes, 1950):

$$VI = (ES/TES) * 100$$

$$O = (SP/TES) * 100$$

$$N = (NP/TP) * 100$$

$$PI = (TPC/TES) * 100$$

Where ES is empty stomach; TES is total examined stomach; SP is number of stomach containing prey; NP is number of each prey item; TP is total number of prey, TPC is total prey category.

Data analysis

All statistics and plots were performed using R version 3.6.1 (R Development Core Team,

<http://www.R-project.org>). The significance level was set at $\alpha = 0.05$ using the function *t-test*.

RESULTS

The averages of physicochemical parameters of Keddara waters Dam are presented in Table 1. Conductivity and pH were significantly higher at the end of the autumn season (December 2019) than at the beginning of the autumn season (September 2020). Unlike the turbidity which was significantly lower at the end of the autumn season (December 2019) than at the beginning of the autumn season (September 2020). The lowest values of transparency were recorded in autumn 2019 (end of the season).

Overall, 67 individuals of Largemouth bass were caught during the autumn of 2019 and 2020. The total length (TL) of all the studied Largemouth bass individuals ranged from 14.2 to 35.6 cm (average length 28.6 ± 0.38 cm for December 2019 and 23.3 ± 2.14 cm for September 2020). Computation of relationships between biometric parameters (TL, FL, and SL) was represented in Fig. 1. All relationships were highly significant (all $R > 0.9$).

The length-weight relationships of Largemouth bass are presented in Fig. 2 and were estimated separately for each year (beginning and end of the autumn season). The sample size and weight ranged from 27.8 cm to 29.8 cm and from 326.16 g to 392.05 g for the end of the autumn season of 2019 and from 14.2 cm to 35.6 cm and from 32.64 g to 722.07 g for the start of the autumn season 2020, respectively. A significant difference was observed between males and females for the length and weight only for the beginning of the autumn season (September 2020).

The values of R^2 and slope b differ from the beginning to the end of the autumn season (December 2019 to September 2020) and all relationships were highly significant, with R^2 values being greater than 0.7. The values of b ranged from 2.29 for the beginning of the autumn season (September 2020), to 3.4 for the end of the autumn season (December 2019).

The total sex ratio of males to females was 2.045:1. The lowest male to female ratio (2:1) was recorded at the start of the autumn season (September 2020) and the highest male to female ratio (2.33:1) was recorded at the end of the autumn season (December 2019) (Table 2).

The composition of *Micropterus salmoides* (female and male) maturity stage for the beginning and the end of the autumn period (Fig. 3) shows that the spawning season starts towards the end of the autumn period for this species at Keddara Dam.

A significant difference was observed ($P < 0.05$) for the condition factor K between beginning and end of the autumn season (December 2019 and September 2020). Largemouth bass reached a higher condition in the end of the autumn season (December 2019) ($K = 15.51 \pm 0.49$) compared to the beginning of the autumn season (September 2020) ($K = 13.92 \pm 2.22$).

For feeding strategy, a total of 60% and 47.4% Largemouth bass contained prey within the stomach for December 2019 and September 2020, respectively, and corresponds to a vacuity index of 20% and 52.6% for the early and late of the autumn season (September 2020 and December 2019), respectively.

Table 3 shows the mean values of diet composition for Largemouth bass. Preys consisted of fish (fish part: 60% and 24.56%) for the beginning and the end of the autumn period (December 2019 and

	pH	Conductivity ($\mu\text{S}/\text{cm}$)	Transparency (m)	Turbidity (NTU)	Salinity (PSU)
December 2019	7.97 ± 0.07	1243 ± 17.6	1.54 ± 0.15	10.01 ± 1.9	0.01
September 2020	7.87 ± 0.05	1039.33 ± 9.77	1.93 ± 0.42	35 ± 3.22	0.01

Table 1. Mean values of the physico-chemical parameters of Keddara waters Dam.




Figure 1. Morphometric relationships between standard, fork, and total length for Black bass caught in Keddara Dam (Algeria). Figure 2. The length-weight relationships for Black bass caught in Keddara Dam (Algeria).

	Male	Female	Total	Ratio
December 2019	7	3	10	2.33:1
September 2020	38	19	57	2:1
Total	45	22	67	2.045:1

Table 2. The sex ratio for male and female *Micropterus salmoides* sampled in Keddara Dam (Algeria).

September 2020), respectively; Full fish: 10% and 26.31% for the beginning and the end of the autumn period (December 2019 and September 2020), respectively), Prawn (10 % and 1.75% for December 2019 and September 2020, respectively), Oligochaeta (Full or Oligochaeta part: 10% and 1.75 % for December 2019 and September 2020, respectively), Insecta (0% and 1.75% for December 2019 and September 2020, respectively). Fish parts constitute the most significant frequently observed prey.

Fish part was more significantly abundant prey in December 2019 than in September 2020 and full fish was the most significantly abundant prey in September 2020 than in December 2019. Largemouth bass exhibits significant predation intensity in December 2019 than in September 2020 for full fish.

DISCUSSION

This research provides information on the biology, presence and ecology of *Micropterus salmoides* in Keddara Dam. Moreover, works on Largemouth bass are few or rare in Algeria despite

Diet	Fish part	Full fish	Prawn	Full Oligochaeta	Oligochaeta part	Insecta
O2019	60*	10	10	10	10	0
O2020	24.56*	26.31	1.75	1.75	1.75	1.75
N2019	55*	5*	10	3.33	6.66	0
N2020	21.66*	22.2*	0.35	0.58	1.17	1.4
PI2019	90	10*	10	10	20	0
PI2020	54.4	47.37*	1.75	1.75	3.51	7.02

Table 3. Diet composition means of Largemouth bass, *Micropterus salmoides* (O: frequency of occurrence (%), N: numerical abundance (%); PI: predation intensity (%), N=10 for December 2019 and N=57 for September 2020; *: significant difference).




Figure 3. Percentages of macroscopic maturity stages in females and males of *Micropterus salmoides*, over the autumn of 2019 and 2020.

its importance in terms of aquaculture and economics. Based on the present results, we can outline an evident adaptation of *Micropterus salmoides* in Keddara Dam that presents excellent possibilities for Largemouth bass growth and reproduction, with favorable environmental conditions.

In fact, it is known that ability of visually feeding piscivorous fish to forage is accepted to be affected by turbidity (Hecht & van der Lingen, 1992). Reid et al. (1999) noted no significant reduction in the diet frequency of Largemouth bass at intermediate levels of turbidity between 18 and 50 NTU, although changes in the prey size selectivity did occur.

Largemouth bass inhabits freshwater environments and sometimes must tolerate sudden fluctuations in salinity. *Micropterus salmoides* appear to have adapted to low-salinity environments by developing osmoregulatory mechanisms, reducing active ion transport, and tolerating higher plasma levels (Meador & Kelso, 1990).

Stroud (1967) reported range of 6.5–8.5 for the optimal pH and Stuber et al., 1982 noted that spawning of Largemouth bass is affected at pH less than 5.0 (no reproduction) and pH above 9.6 (no eggs survival).

In aquatic communities, light is an important factor that influences predator-prey interactions (McMahon & Holanov, 1995). Baker et al. (1993) reported that Largemouth bass seek protection from light and prefer shaded areas during all stages of life. McMahon & Holanov (1995) indicated that water clarities less than 0.5 m Secchi depth reduce Largemouth bass foraging success to 67 and 75% less than at 2 m Secchi depth and high water clarities (4 m Secchi depth).

Regarding conductivity, Glover et al. (2012, 2013) and Huntsman et al. (2021) reported water quality conditions within Largemouth bass physiological limits of conductivity $< 8600 \mu\text{s}\cdot\text{cm}^{-1}$.

In this study, the observed differences in length-weight relationships can be related to reproduction or feeding activities (Weatherley & Gill, 1987; Wootton, 1990; Moutopoulos & Stergiou, 2002), and the differences in b-values can be attributed to the combination of one or more factors such as differences in the number of individual examined, differences in the observed length ranges of the fish caught and season effect (Moutopoulos & Stergiou, 2002, Chabet dis et al., 2022).

Because of the size-selective properties of the trammel net used, the samples do not include the small-sized individuals for Largemouth bass caught. Consequently, the use of the length-weight relationships in the present study should be limited to the observed length ranges (Petrakis & Stergiou, 1995; Goncalves et al., 1996; Chabet dis et al., 2022).

The sexual dimorphism in the size of *Micropterus salmoides* was reported in the first studies published by Heidinger (1976). Also, larger females were observed by Alessio (1984) only occasionally. However, Rodríguez-Sánchez et al. (2009) noted no significant differences between sexes in Primera de Palos' lake (Huelva, Spain). In Keddara Dam, no significant differences were detected between males and females at the end of the autumn season (December 2019) unlike at the start of the autumn season (September 2020) when a significant difference was observed between males and females for the length and weight. These results support the results of other researchers and can be explained by reproductive activity (Beamish et al., 2005; Granado-Lorencio, 2000).

The sex ratio of males to females of *Micropterus salmoides* from Keddara Dam was 2.33 males: 1 female for December 2019 and 2.33 males: 1 female for September 2020. This result seems close to the result of Beamish et al. (2005) in Lake Manyame, Zimbabwe (2.05 males: 1 female), and different from the values found by Taylor & Weyl (2017) in Wriggleswade (1 male: 1.13 females) and in Mankazana (1 male: 1.14 females) in the Eastern Cape Province of South Africa. A number of hypotheses may explain these subsequent changes in this value, like differences in habitat preference according to the sex or season, selective mortality, sampling errors, reproductive stress suffered by female and males (Fernandez & Rossomanno, 1997; Granado-Lorencio, 2000; Beamish et al., 2005).

Regarding the diet of Largemouth bass *Micropterus salmoides* in the Algerian freshwaters, there is very scarce data. The present finding showed that Largemouth bass is not exclusively piscivorous, feeds also on prawns, Oligochaeta, and insects that live in Keddara Dam. Marinelli et al., (2007) mentioned that decapod *Palaemonetes antennarius* is the major food items *Micropterus salmoides* in Lake Bracciano (Italy). Alessio (1984), Rodriguez-Jimenez (1989), Godinho & Fer-

reira (1994), Nicola et al. (1996), Godinho et al. (1997) and Godinho & Ferreira (1998) indicated that Largemouth bass ingest fish (the predominant food item) and crustaceans and insects (also frequently eaten).

The number of prey was lower in our study when compared with other studies (Rodríguez-Jimenez, 1989; Godinho & Ferreira, 1994; Godinho et al., 1997; Godinho & Ferreira, 1998; Marinelli et al., 2007). He et al. (1994), Hickley et al. (1994), Godinho et al. (1997) and Olson et al. (1998) reported that Black bass is an opportunistic predator with an alimentary range depending on the feeding resources and on the environment.

The current study showed a significant difference related to the stomach content of fish (fish part or full). Many factors could have affected the amount and type of food found in the stomach content of Black bass like the water temperature and the reduction of the foraging activity during the reproductive period (Alessio, 1984; Roseblum et al., 1994).

The feeding and spawning strategy in Keddara Dam is likely an adaptation for Largemouth bass that helps ensure reproductive and life success where the biotope is frequently subjected to disturbances out of control, like water level fluctuations, noting that these waters are intended for drinking water. However, in freshwater environments, biological invasions are an important factor for biodiversity loss (Almeida et al., 2012) and have a variety of negative impacts on habitats and native species, such as habitat degradation (García-Berthou, 2001), competition (Keller & Brown, 2008), food web alteration (Almeida et al., 2009), and predation (Schilling et al., 2009).

In conclusion, the present study provides some basic ecologic and biologic information on the reproduction and feeding habits of *Micropterus salmoides* in Algeria, reporting for the first time the presence of this species in a favourable environment as Keddara Dam.

REFERENCES

- Alessio G., 1984. Le black-bass (*Micropterus salmoides* Lacép.) dans les eaux italiennes. Un antagoniste du brochet? Bulletin Français de Pisciculture, 292: 1–17.
- Almeida D., Almodóvar A., Nicola G.G. & Elvira B., 2009. Feeding tactics and body condition of two introduced populations of pumpkinseed *Lepomis gibbosus*: taking advantages of human disturbances?. Ecology of freshwater fish, 18: 15–23.
- Almeida D., Almodóvar A., Nicola G.G., Elvira B. & Grossman G.D., 2012. Trophic plasticity of invasive juvenile largemouth bass *Micropterus salmoides* in Iberian streams. Fisheries Research, 113: 153–158.
- Azuma M. & Motomura Y., 1998. Feeding habits of Largemouth bass in a non-native environment: The case of a small lake with bluegill in Japan. Environmental Biology of Fishes, 52: 379–389.
- Bai J. & Li S., 2019. Genetic Breeding and Molecular Marker-Assisted Selective Breeding of Largemouth Bass, 1st ed. Academic Press by Elsevier Inc., 332 pp. <https://doi.org/10.1016/C2017-0-04751-1>
- Baker J.P., Warren-Hicks W.J., Gallagher J. & Christensen S.W., 1993. Fish population losses from Adirondack lakes: the role of surface water acidity and acidification. Water Resources Research, 29: 861–874.
- Beamish C.A., Booth A.J. & Deacon N., 2005. Age, growth and reproduction of largemouth bass, *Micropterus salmoides*, in lake Manyame, Zimbabwe. African Zoology, 40: 63–69.
- Britton J.R., Harper D.M. & Oyugi D.O., 2010. Is the fast growth of an equatorial *Micropterus salmoides* population explained by high water temperature?. Ecology of Freshwater Fish, 19: 228–238.
- Cai J., Zhou X., Yan X., Lucente D. & Lagana C., 2019. Top 10 species groups in global aquaculture 2017. WAPI Factsheet. FAO, 12 pp.
- CFS, 2019. China Fishery Statistics Yearbook. Bureau of Fisheries, Ministry of Agriculture, China Agriculture Press.
- Chabet dis C., Kara-Toumi F.Z., Bouamama S., Sahra F. & Boucena M.A., 2022. Eco-biological Study of the Mosquitofish *Gambusia affinis* from Oubeira Lake. Journal of Aquaculture and Fish Health, 11: 201–209. <https://doi.org/10.20473/jafh.v11i2.30017>
- Cooke S. & Philipp D.P., 2009. Centrarchid Fishes: Diversity, Biology and Conservation. Wiley-Blackwell, 560 pp. <https://doi.org/10.1002/9781444316032>
- Fernández-Delgado C. & Rossomanno S., 1997. Reproductive biology of the mosquitofish in a permanent natural lagoon in south-west Spain: two tactics for one species. Journal of Fish Biology, 51: 80–92.
- García-Berthou E., 2001. Size- and depth-dependent variation in habitat and diet of the common carp (*Cyprinus carpio*). Aquatic Sciences, 63: 466–476.
- Glover D.C., DeVries D.R. & Wright R.A., 2012. Effects of temperature, salinity and body size on routine metabolism of coastal Largemouth bass

- Micropterus salmoides*. Journal of Fish Biology, 81: 1463–1478.
<https://doi.org/10.1111/j.1095-8649.2012.03385.x>
- Glover D. C., DeVries D.R. & Wright R.A., 2013. Growth of Largemouth bass in a dynamic estuarine environment: an evaluation of the relative effects of salinity, diet, and temperature. Canadian Journal of Fisheries and Aquatic Sciences, 70: 485–501.
- Godinho F.N. & Ferreira M.T., 1994. Diet composition of Largemouth black bass, *Micropterus salmoides* (Lacepede), in southern Portuguese reservoirs: its relation to habitat characteristics. Fisheries Management and Ecology, 1: 129–137.
- Godinho F.N. & Ferreira M.T., 1998. Spatial variation in diet composition of pumpkinseed sunfish, *Lepomis gibbosus*, and Largemouth bass, *Micropterus salmoides*, from a Portuguese stream. Folia Zoologica-Praha, 47: 205–214.
- Godinho F., Ferreira M.T. & Cortes R.V., 1997. The environmental basis of diet variation in pumpkinseed sunfish, *Lepomis gibbosus*, and Largemouth bass, *Micropterus salmoides*, along an Iberian river basin. Environmental Biology of Fishes, 50: 105–115.
- Gonçalves J.M.S., Bentes L., Lino P.G., Ribeiro J., Canario A.V. & Erzini K., 1996. Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and south-west coast of Portugal. Fisheries Research, 30: 253–256.
- Granado-Lorencio C., 2000. Ecología de comunidades. El paradigma de los peces de agua dulce. University of Seville, Spain, 282 pp.
- He X., Hodgson J. R., Kitchell J.F. & Wright R.A., 1994. Growth and diet composition of Largemouth bass (*Micropterus salmoides*) from four experimental lakes. Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen, 25: 2092–2097.
- Hecht T. & Van der Lingen C.D., 1992. Turbidity-induced changes in feeding strategies of fish in estuaries. African Zoology, 27: 95–107.
- Heidinger R.C., 1976. Synopsis of biological data on the Largemouth bass *Micropterus salmoides* (Lacepede, 1802). FAO Fisheries Synopses, 115: 1–85.
- Hickley P., North R., Muchiri S.M. & Harper D.M., 1994. The diet of Largemouth bass, *Micropterus salmoides*, in Lake Naivasha, Kenya. Journal of Fish Biology, 44: 607–619.
- Holden M.J. & Raitt D.F.S., 1974. Manuel des sciences halieutiques. Deuxième partie. Méthodes et recherches sur les ressources et leur application. Rome, Italy, FAO, Document Technique sur les Pêches, (115), Rev. 1., 223 pp.
- Huntsman B.M., Feyrer F., Young M. J., Hobbs J.A., Acuña S., Kirsch J. E., Mahardja B. & Teh S., 2021. Recruitment dynamics of non-native Largemouth bass within the Sacramento–San Joaquin Delta. Canadian Journal of Fisheries and Aquatic Sciences, 78: 505–521.
- Hussein G.H.G., Chen M., Qi P.P., Cui Q.K., Yu Y., Hu W.H., Tiana Y., Fan Q.X., Gao Z.X., Feng M.W. & Shen Z.G., 2019. Aquaculture industry development, annual price analysis and out-of-season spawning in Largemouth bass *Micropterus salmoides*. Aquaculture, 734901.
<https://doi.org/10.1016/j.aquaculture.2019>
- Hynes H.B.N., 1950. The food of fresh-water sticklebacks (*Gasterosteus aculeatus* and *Pygosteus pungitius*), with a review of methods used in studies of the food of fishes. The Journal of Animal Ecology, 19: 36–58.
- Kara H.M., 2012. Freshwater fish diversity in Algeria with emphasis on alien species. European journal of wildlife research, 58: 243–253
- Keller K. & Brown C., 2008. Behavioural interactions between the introduced plague minnow *Gambusia holbrooki* and the vulnerable native Australian ornate rainbowfish *Rhadinocentrus ornatus*, under experimental conditions. Journal of Fish Biology, 73: 1714–1729.
- Marinelli A., Scalici M., & Gibertini G., 2007. Diet and reproduction of Largemouth bass in a recently introduced population, Lake Bracciano (Central Italy). Bulletin Français de la Pêche et de la Pisciculture (385): 53–68.
- McLean E., Alfrey K.B., Gatlin III D.M. & Barrows F.T., 2022. Responses of largemouth bass (*Micropterus salmoides*, Lacépède, 1802) to fishmeal-, and fish oil-free diets. Aquaculture Research, 53: 3036–3047.
<https://doi.org/10.1111/are.15815>
- McMahon T.E. & Holanov S.H., 1995. Foraging success of Largemouth bass at different light intensities: implications for time and depth of feeding. Journal of Fish Biology, 46: 759–767.
- Meador M.R. & Kelso W.E., 1990. Physiological responses of Largemouth bass, *Micropterus salmoides*, exposed to salinity. Canadian Journal of Fisheries and Aquatic Sciences, 47: 2358–2363.
<https://doi.org/10.1139/f90-262>
- Moutopoulos D.K. & Stergiou K.I., 2002. Length–weight and length–length relationships of fish species from the Aegean Sea (Greece). Journal of Applied Ichthyology, 18: 200–203.
- Nicola G.G., Almodovar A. & Elvira B., 1996. The diet of introduced Largemouth bass, *Micropterus salmoides*, in the Natural Park of the Ruidera Lakes, central Spain. Polskie Archiwum Hydrobiologii, 43: 2.
- Olson M.H., Carpenter S.R., Cunningham P., Gafny S., Herwig B.R., Nibbelink N.P., Pellet T., Storlie C.,

- Trebitz A.S. & Wilson K.A., 1998. Managing macrophytes to improve fish growth: a multi-lake experiment. *Fisheries*, 23: 6–12.
- Petrakis G. & Stergiou K.I., 1995. Weight-length relationships for 33 fish species in Greek waters. *Fisheries research*, 21: 465–469.
- Reid S.M., Fox M.G., & Whillans T.H., 1999. Influence of turbidity on piscivory in Largemouth bass (*Micropterus salmoides*). *Canadian Journal of Fisheries and Aquatic Sciences*, 56: 1362–1369.
- Rodriguez-Jimenez A.J., 1989. Hábitos alimenticios de *Micropterus salmoides* (Pisces: Centrarchidae) *Lepomis gibbosus* (Pisces: Centrarchidae) y *Gambusia affinis* (Pisces: Poeciliidae) en las orillas del embalse de prosperina (Extremadura, España). *Limnética*, 5: 13–20.
- Rodríguez-Sánchez V., Encina L., Rodríguez-Ruiz A. & Sánchez Carmona R., 2009. Largemouth bass, *Micropterus salmoides*, growth and reproduction in Primera de Palos' lake (Huelva, Spain). *Folia Zoologica Praha*, 58: 436–446.
- Rosenblum P.M., Brandt T.M., Mayes K.B. & Hutson P., 1994. Annual cycles of growth and reproduction in hatchery-reared Florida Largemouth bass, *Micropterus salmoides floridanus*, raised on forage or pelleted diets. *Journal of Fish Biology*, 44: 1045–1059.
- Schilling E.G., Loftin C.S. & Huryn A.D., 2009. Effects of introduced fish on macroinvertebrate communities in historically fishless headwater and kettle lakes. *Biological Conservation*, 142: 3030–3038.
- Stroud R.H., 1967. Water quality criteria to protect aquatic life: a summary. American Fisheries Society Special Publication, 4: 33–37.
- Stuber R.J., Gebhart G. & Maughan O.E., 1982. Habitat suitability index models: Largemouth bass. Western Energy and Land Use Team, Office of Biological Services, Fish and Wildlife Service, US Department of the Interior. FWS/OBS-82/10.16, 32 pp.
- Taylor G.C. & Weyl O.L.F., 2017. Age, growth and reproduction of non-native Largemouth bass *Micropterus salmoides* (Lacépède, 1802) populations in two temperate African impoundments. *Journal of Applied Ichthyology*, 33: 767–775. <https://doi.org/10.1111/jai.13353>
- Weatherley A.H. & Gill H.S., 1987. The biology of fish growth. Academic Press, London. 443 pp.
- Wootton R.J., 2012. Ecology of teleost fish. Chapman & Hall, London, 756 pp.
- Zouakh D.E. & Meddour A., 2017. First experimental induced breeding of the Largemouth Bass *Micropterus salmoides* Lacépède, 1802 (Centrarchidae) in Algeria. *Journal of Applied Environmental and Biological Sciences*, 7: 1–10.