

## A new species of *Alvania* Risso, 1826 from the Western Indian Ocean (Gastropoda Rissoidae)

Bruno Amati

<sup>1</sup>Largo Giuseppe Veratti, 37/D, 00146 Roma, Italy; e-mail: bruno\_amati@yahoo.it, <https://orcid.org/0000-0002-3515-9543>

---

### ABSTRACT

A new species of the rissoid genus *Alvania* Risso, 1826 (Gastropoda Rissoidae) is described based on shells sorted out of a bioclastic sand sample collected with a trap box for *Marginella* species in Manda Toto Island (Lamu Archipelago, Kenya). It is compared with the most similar congeners.

### KEY WORDS

Gastropoda; Rissoidae; Western Indian Ocean; *Alvania*; new species; taxonomy.

Received 21.12.2023; accepted 18.01.2024; published online 08.02.2024

---

### INTRODUCTION

The genus *Alvania* Risso, 1826 (Gastropoda Rissoidae) as currently conceived includes a speciose group of small to large sized rissoid species living from the lower intertidal to bathyal depths, mostly on the continental shelf, where they are prevalently associated with algal facies (e.g.: Ponder, 1985; Gofas, 2007; Ávila et al., 2012), with few representatives in deeper waters down to 4700 m depth (Bouchet & Warén, 1993). They feed on diatoms, dinoflagellates, and detritus, with the deep-sea species presumably being selective deposit feeders (Ponder, 1985).

The genus *Alvania* comprises 279 known Recent species and 201 fossil ones (MolluscaBase, 2024) from the northeastern Atlantic Ocean and Mediterranean Sea, the northwestern Atlantic and the Caribbean, the eastern Pacific Ocean, the Indo-West Pacific Ocean, the temperate Australia, and South Africa (Ponder, 1985).

The Rissoidae are not considered in the limited general literature concerning the malacofauna of

Kenya (e.g.: Fowler, 2016). Furthermore, the few authors that have worked on rissoids from Kenya, have not, or only marginally treated the genus *Alvania* (e.g.: Bosch et al., 1998; Callea et al., 2005; Amati, 2024). A new species has been found in a bioclastic sand sample for Kenya, and it is described here

### MATERIAL AND METHODS

The specimens studied herein have been sorted out a bioclastic sand sample, collected with a trap box for *Marginella* species, in Manda Toto Island (Lamu Arcipelago, Kenya) -2.21817, 41.01517, at approximately 2 m depth, in February 2023. The shells were studied and measured under a Kyowa stereomicroscope, with a micrometric eyepiece (with magnification X90, ± 10% deviation), using the method of Verduin (1982). Photographs have been taken with a Sony Cyber-shot DSC-W110 digital camera mounted on a Kyowa KBS and a Kyowa SDZ-P stereomicroscopes, edited with the software Combine-Z (Hadley, 2006). Current systematics is based on MolluscaBase (2024).

**ABBREVIATIONS AND ACRONYMS.** AMS: The Australian Museum (Sydney, Australia); BA: Bruno Amati collection (Roma, Italy); dd: empty shell(s). MNHN: National Museum of Natural History (Paris, France); ZMB: Museum für Naturkunde – Leibniz-Institut für Evolutions- und Biodiversitätsforschung an der Humboldt-Universität zu Berlin, Berlin (Germany); vs: *versus*.

## RESULTS

### Systematics

Classis GASTROPODA Cuvier, 1795  
 Subclassis CAENOGASTROPODA Cox, 1960  
 Ordo LITTORINIMORPHA Golikov et Starobogatov, 1975  
 Superfamilia RISSOOIDEA Gray, 1847  
 Familia RISSOIDAE Gray, 1847  
 Genus *Alvania* Risso, 1826

**TYPE SPECIES.** *Alvania europea* Risso, 1826: 142, pl. IX, fig. 116 synonym of *Alvania cimex* (Linnaeus, 1758) by subsequent designation (Nevill, 1885: 105).

***Alvania pettinellii* n. sp.** (Figs 1; 2–8; Table 1)  
<https://www.zoobank.org/A67F5EB7-588B-4BDF-ACF4-7804BD83B3A1>

**DIAGNOSIS.** *Alvania* small for the genus (height <2 mm), with rather robust shell; protoconch paucispiral; teleoconch morphologically scarcely variable; spiral sculpture predominant over the axial one. Coloration yellowish with quadrangular brownish subsutural and basal blotches.

**TYPE MATERIAL.** Holotype. KENYA • dd (height 1.57 mm, width 0.92 mm, Figs 2–5, 7, 8); Lamu Archipelago, Manda Toto Island; -2.21817, 41.01517; 2 m; bioclastic sand; Feb. 2023; Domenico Pettinelli legit; MNHN-IM-2000-39447. Paratypes. KENYA • 2 dd; same locality data as holotype; BA.


**TYPE LOCALITY.** Manda Toto Island, Lamu Archipelago, Kenya; 2.21817, 41.01517; 2 m depth.

**DESCRIPTION OF THE HOLOTYPE.** Shell (Figs 2–4). Small for the genus, height 1.57 mm, width 0.92 mm, height/width ratio 1.71, solid, ovate-conical. Protoconch (Figs 5, 7, 8) paucispiral, dome-shaped, of 1.3 whorls, height 0.276 mm, nucleus diameter 0.113 mm, first half whorl diameter 0.188 mm, maximum diameter 0.363 mm; sculptured by 9 thin cordlets, slightly wrinkled. Protoconch-teleoconch boundary well marked, slightly opisthocline.

Teleoconch of 3.1 convex whorls, with sutures impressed. Spiral sculpture stronger than axial, of non-equidistant smooth cords, 8 on the last whorl, 3 above the aperture and 5 on the base. Cords I, II



Figure 1. Type locality (red dot) of *Alvania pettinellii* n. sp., Indian Ocean, Kenya, Lamu Archipelago, Manda Toto Island.



Figures 2–8. *Alvania pettinellii* n. sp. Figs 2–5, 7, 8: holotype, Manda Toto Island (Kenya), height 1.57 mm (MNHN-IM-2000-39447). Fig. 6: paratype, Manda Toto Island (Kenya), height 1.73 mm (BA). Scale images 2–4, 6. Scale bars: 5, 7, 8, 100 µm.

and III starting immediately after the protoconch-teleoconch boundary (Fig. 5); interspace between cords I and II slightly larger than between cords II and III. Numerous thin orthoclin3 axial riblets in the interspaces, c. 35 on the last whorl, interrupted on spiral cord IV; no thickening at the intersections. Umbilical fissure absent. Aperture pyriform, height 0.67 mm, height/aperture height ratio 2.34, peristome continuous, outer lip sharp, internally smooth, varix broad, slightly prosocline.

Colour: teleoconch yellowish with quadrangular brownish subsutural and basal blotches, and a darker subsutural spot on the outer lip. Initial part of the protoconch (nucleus and first half whorl) dark brown (Fig. 8).

Operculum and soft parts unknown.

**VARIABILITY.** Only three specimens examined, two adults and one juvenile: no significant variation detected (Table 1). Maximum measurements: height 1.73 mm, width 0.97 mm, height/width ratio 1.78, aperture height 0.67 mm, height/ aperture height ratio 2.58 (Fig. 6).

**ETYMOLOGY.** Named after Domenico Pettinelli (Roma) who collected the bioclastic sand sample in which the new species was found and made it available for study.

**DISTRIBUTION.** *Alvania pettinellii* n. sp. so far known only on empty shells from the type locality: the western Indian Ocean in East Africa (Kenya), in 2 m depth (Fig. 1).

**REMARKS.** *Alvania isolata* (Laseron, 1956) from Australia (see Laseron, 1956: 439, fig. 144; Ponder, 1985: 142, figs 93 A, B; Amati et al., 2023: 808, figs 4A–H; 48; Fig. 9), differs from *A. pantanellii* n. sp.

in its teleoconch with axial and spiral sculpture of equal thickness; the four spiral cords on the whorls vs three in *A. pettinellii* n. sp.; the protoconch with 4–5 smooth spiral cordlets vs 9 thin cordlets in *A. pettinellii* n. sp.; the different chromatic pattern (whitish, with series of pairs of short brown lines vs yellowish with quadrangular brownish subsutural and basal blotches in *A. pettinellii* n. sp.).

*Alvania albachiarae* Perugia, 2021 from Oman (see Perugia, 2021: 1, pl. 1, figs. d–f), differs from *A. pettinellii* n. sp. in its four spiral cords on the teleoconch vs three in *A. pettinellii* n. sp.; the cords I, II, III and IV starting immediately after the protoconch-teleoconch boundary vs cords I, II, III in *A. pettinellii* n. sp.; the axials thicker and fewer, forming tubercles at the intersections with spirals.

*Alvania denseclathrata* Thiele, 1925 from southern Africa, (see Thiele, 1925: 81, pl. 6, fig. 11; Amati et al., 2023: 806, figs 2A–G; 48; Fig. 10), differs from *A. pettinellii* n. sp. in its larger size (height 2.2 mm vs 1.73 mm in *A. pettinellii* n. sp.); the axial and spiral sculpture denser, of equal thickness vs spirals thicker than axials in *A. pettinellii* n. sp.; the coloration uniform whitish, devoid of spots, present in *A. pettinellii* n. sp.

*Alvania herosae* Amati, Di Giulio et Oliverio, 2024 from French Polynesia (see Amati et al., 2023: 811, figs 7A–E; 8A–E; 9A–F; 10A–D; 11A–D; 12A–E; 13A–I; 49A; 53B, C; Fig. 11), differs from *A. pettinellii* n. sp. in its more robust axial sculpture which, at the intersection with the spiral cords, forms evident tubercles, absent in *A. pettinellii* n. sp.; the different chromatic pattern (protoconch monochrome and columellar area brownish vs nucleus and first half whorl dark brown, and columellar area yellowish in *A. pettinellii* n. sp.).

Protoconch	1	2	3	Min–max	Mean	St. dev.
Height	0.276	0.263	0.238	0.238–0.276	0.259	0.019
Diameter of the nucleus	0.113	0.125	0.113	0.113–0.125	0.117	0.007
Diameter first half whorl	0.188	0.225	0.213	0.188–0.225	0.209	0.019
Maximum diameter	0.363	0.350	0.325	0.325–0.363	0.346	0.019
N° of whorls	1.3	1.3	1.3	1.3	1.3	0
Paucispiral	yes	yes	yes	yes	yes	

Table 1. Measurements of the protoconch of *Alvania pettinellii* n. sp. in mm, with range, mean, and standard deviation. 1. Holotype, 2, 3 paratypes. All from the type locality.




Figure 9. *Alvania isolata* (Laseron, 1956), holotype, height 1.5 mm, Christmas Island, depth unknown, AMS C.102476, (photo: A.C. Miller; copyright: Australian Museum). Figure 10. *Alvania denseclathrata* (Thiele, 1925), lectotype, height 2.2 mm, South Africa, Valdivia Stn 95, 80 m, ZMB/Moll. N° 64984. Figure 11. *Alvania herosae* Amati, Di Giulio et Oliverio, 2023, holotype, Australes, Rapa, Tupuaki Bay, Atelier RAPA Stn 21, 5 m, height 2.02 mm, MNHN-IM-2000-38704. Figure 12. *Alvania uapou* Amati, Di Giulio et Oliverio, 2023, holotype, height 1.70 mm, Marquesas, Ua Pou, Motu Mokohe, Atelier MARQUISES Stn 20, 10–15 m, MNHN-IM-2000-38710. Scale images.

*Alvania uapou* Amati, Di Giulio et Oliverio, 2024 from French Polynesia (see Amati et al., 2023: 824, figs 18A–F; 19A–E; 49C; 53F; Fig. 12) differs from *A. pettinellii* n. sp. in its most robust and fewer axials; the protoconch sculptured with seven zig-zag spiral threads vs nine thin cordlets in *A. pettinellii* n. sp.

*Alvania* sp. 1 (see Hasekawa, 2006: 107, 108, fig. 3) from Philippines differs from *A. pettinellii* n. sp. in its spiral and axial sculpture of equal robustness vs spirals thicker than axials in *A. pettinellii* n. sp.; the aperture internally lyrate vs smooth in *A. pettinellii* n. sp.; the different chromatic pattern (with small brownish spots scattered on the whitish background vs yellowish with quadrangular brown subsutural and basal blotches in *A. pettinellii* n. sp.).

## ACKNOWLEDGEMENTS

I am grateful to Domenico Pettinelli (Roma, Italy) for presenting to me the bioclastic sand sample in which the new species was found, and to Marco

Oliverio (Department of Biology & Biotechnologies ‘Charles Darwin’, Sapienza University of Rome, Italy) for his very helpful comments. I thank Christine Zorn (ZMB) and Alison Miller, Mandy Reid and Francesco Criscione (AMS), who helped with type material in the collections under their care; Virginie Héros (MNHN, Paris, France) and Roberto Ardonini (Roma, Italy) for bibliographical help.

## REFERENCES

- Amati B., 2024. A new species of *Merelina* Iredale, 1915 from the Western Indian Ocean (Caenogastropoda: Lironobidae). *Bollettino Malacologico*. In press.
- Amati B., Di Giulio A. & Oliverio M., 2023. Shallow-water Rissoidae of the genera *Alvania* Risso, 1826, *Haurakia* Iredale, 1915, *Parashiela* Laseron, 1956, *Simulamerelina* Ponder, 1985 and *Subestea* Cotton, 1944 (Gastropoda, Caenogastropoda, Rissooidea) from French Polynesia, with the description of a new deep-water genus. *Zoosystema*, 45: 803–892.

- [https://doi.org/10.5252/zosistema2023v45a25.](https://doi.org/10.5252/zosistema2023v45a25)  
<http://zosistema.com/45/25>
- Ávila S. P., Goud J. & De Frias Martins A. M., 2012. Patterns of diversity of the Rissoidae (Mollusca: Gastropoda) in the Atlantic and the Mediterranean region. *The Scientific World Journal*, no. 164890.  
<https://doi.org/10.1100/2012/164890>
- Bartsch P., 1915. Report on the Turton Collection of South African marine mollusks, with additional notes on other South African shells contained in the United States National Museum. *Bulletin. United States National Museum*. 91: xii + 305 pp. + 54 plates.
- Bosch, D. T., Dance S. P., Moolenbeek R. G. & Oliver P. G., 1995. Seashells of eastern Arabia. Dubai: Motivate publishing. Edit by S. Peter Dance, 296 pp.  
<https://hdl.handle.net/11245/1.102842>
- Bouchet P. & Warén A., 1993. Revision of the Northeast Atlantic bathyal and abyssal Mesogastropoda. *Bollettino Malacologico*, Supplemento 3: 579–849.  
<https://doi.org/10.5962/bhl.title.140732>
- Bozzetti L., 2017. *Alvania hueti* (Gastropoda: Hortugastropoda: Rissoidae) nuova specie dall'isola di Reunion. *Malacologia Mostra Mondiale*, 96: 6–7.
- Callea A., Borri M., Cianfanelli S., Martignoni R. & Volpi C., 2005. Taxonomic and ecological remarks on the marine molluses of the Mida Creek area (Kenya, Western Indian Ocean). *Atti Società italiana Scienze naturali Museo civico Storia naturale Milano*, 146: 74–94.
- Cecalupo A. & Perugia I., 2009. New species from South Madagascar (2<sup>nd</sup> note). *Malacologia Mostra Mondiale* 63: 21.
- Fowler O., 2016. Seashells of the Kenya coast. Conc Books: Harxheim. 170 pp.
- Gofas S., 2007. Rissoidae (Mollusca: Gastropoda) from northeast Atlantic seamounts. *Journal of Natural History*, 41: 779–885.
- Hadley A., 2006. Combine ZP public domain image processing software. Available from <https://web.archive.org/web/20160221032141/http://www.hadleyweb.pwp.blueyonder.co.uk/>
- Hasegawa K., 2006. A small collection of rissoid gastropods (Mollusca: Gastropoda: Rissoidae) collected from Puerto Galera, Mindoro Island, the Philippines.
- Memoirs of the National Science Museum, Tokyo (44): 105–118.
- Krauss F., 1848. Die Südafrikanischen Mollusken. Ein Beitrag zur Kenntniss der Mollusken des Kap- und Natallandes und zur Geographischen Verbreitung derselben mit Beschreibung und Abbildung der neuen Arten. Ebner and Seubert, Stuttgart, 140 pp., 6 pls.
- Laseron C.F., 1956. The Families Rissoinidae and Rissoidae (Mollusca) from the Solanderian and Dampierian zoogeographical Provinces. *Australian Journal of Marine and Freshwater Research*, 7: 384–484.
- Melvill J.C. & Standen R., 1901. The Mollusca of the Persian Gulf, Gulf of Oman and Arabian Seas as evidenced mainly through the collections of Mr. F. W. Townsend, 1893–1900, with descriptions of new species. Part 1, Cephalopoda, Gastropoda, Scaphopoda. *Proceedings of the Zoological Society of London*, 1901 (2): 327–460.
- MolluscaBase Eds, 2024. *Alvania* Risso, 1826. Accessed at <https://www.molluscabase.org/aphia.php?p=taxdetails&id=138439> on 2 March 2023.
- Nevill G., 1885. Hand list of Mollusca in the Indian Museum, Calcutta. Part II. Gastropoda, Prosobranchia–Neurobranchia (contd.). Government Printer, Calcutta, 306 pp.
- Perugia I., 2021. Some new microshells from Oman (Masirah I.). *Xenophora Taxonomy*, 32: 1–9.
- Ponder W. F., 1985. A review of the genera of the Rissoidae (Mollusca: Mesogastropoda: Rissoacea). *Records of the Australian Museum*, Supplement 4: 1–221.
- Risso A., 1826. Des principales productions de l'Europe Méridionale. Vol. 4. Histoire naturelle, Paris, 439 pp., 12 pls.
- Thiele J., 1925. Gastropoden der Deutschen Tiefsee-Expedition. II Teil. Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898–1899: 17 (2): 35–382, pls 13–46 [reprints paginated 1–348, pls 1–34].
- Verduin A., 1982. How complete are diagnoses of coiled shells of regular build? A mathematical approach. *Basteria*, 45: 127–142.